

Firman Alamsyah

Non-contact electric field exposure provides potential cancer therapy through p53-independent proliferation arrest and intri...

-  Biologi
-  Permintaan Dosen (2026)
-  Universitas Al Azhar Indonesia

Document Details

Submission ID

trn:oid:::1:3475021536

10 Pages

Submission Date

Feb 7, 2026, 9:19 AM GMT+7

6,389 Words

Download Date

Feb 7, 2026, 10:40 AM GMT+7

34,312 Characters

File Name

1._Non-contact_electric_fields_exposure.pdf

File Size

1.1 MB

15% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- ▶ Bibliography

Exclusions

- ▶ 9 Excluded Sources
- ▶ 33 Excluded Matches

Match Groups

-  **32** Not Cited or Quoted 8%
Matches with neither in-text citation nor quotation marks
-  **31** Missing Quotations 8%
Matches that are still very similar to source material
-  **0** Missing Citation 0%
Matches that have quotation marks, but no in-text citation
-  **0** Cited and Quoted 0%
Matches with in-text citation present, but no quotation marks

Top Sources

- 11%  Internet sources
- 12%  Publications
- 3%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

-  **32** Not Cited or Quoted 8%
Matches with neither in-text citation nor quotation marks
-  **31** Missing Quotations 8%
Matches that are still very similar to source material
-  **0** Missing Citation 0%
Matches that have quotation marks, but no in-text citation
-  **0** Cited and Quoted 0%
Matches with in-text citation present, but no quotation marks

Top Sources

- 11%  Internet sources
- 12%  Publications
- 3%  Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

Rank	Type	Source	Percentage
1	Internet	www.ncbi.nlm.nih.gov	2%
2	Internet	jurnalpenyuluhan.ipb.ac.id	2%
3	Internet	f1000research.com	1%
4	Student papers	Institut Pertanian Bogor	<1%
5	Internet	aacrjournals.org	<1%
6	Internet	www.frontiersin.org	<1%
7	Internet	open.library.ubc.ca	<1%
8	Internet	ejournal.undip.ac.id	<1%
9	Student papers	Oxford Brookes University	<1%
10	Internet	karger.com	<1%

11	Internet	
	www.researchgate.net	<1%
12	Internet	
	www.mdpi.com	<1%
13	Publication	
	Sahudi Abdul Mujib, Firman Alamsyah, Warsito Purwo Taruno. "Cell Death and In...	<1%
14	Internet	
	www.spandidos-publications.com	<1%
15	Publication	
	Vinod Kumar Gurjar, Shweta Jain, Ankur Vaidya, Kuldeep Bansal. "Reinstating the ...	<1%
16	Internet	
	www.nature.com	<1%
17	Internet	
	link.springer.com	<1%
18	Publication	
	Debajani Mohanty, Sucheesmita Padhee, Arpita Priyadarshini, Bibhuti Bhushan Ch...	<1%
19	Publication	
	Dwiyati Pujimulyani, Ch. Lilis Suryani, Astuti Setyowati, Rr. Anisa Siwianti Handay...	<1%
20	Publication	
	Maryam Khezri, Gholamreza Salehi Jouzani, Masoud Ahmadzadeh. "Fusarium cul...	<1%
21	Publication	
	Role of Proteases in Cellular Dysfunction, 2014.	<1%
22	Publication	
	S W Purnami, R S Putra, A I Edina, I N Pertiwi, E Sukur, N Soraya. "Cox Model Surv...	<1%
23	Publication	
	Wahyu Widowati, Lusiana Darsono, Johan Lucianus, Edwin Setiabudi et al. "Butter...	<1%
24	Internet	
	d-nb.info	<1%

25 Internet
mdpi-res.com <1%

26 Publication
"Novel Apoptotic Regulators in Carcinogenesis", Springer Science and Business M... <1%

27 Publication
Ebru Emekli-Alturfan. "Zebrafish Models of Neurodegenerative Disorders", CRC Pr... <1%

28 Publication
Gaia Palmini, Francesca Marini, Maria Brandi. "What Is New in the miRNA World R... <1%

29 Publication
Silveira, Aleth  ia L., Gla  cia V. Faheina-Martins, Raquel C. Maia, and Demetrius A. ... <1%

30 Publication
Ying Li, Hai-jun Xiao, Feng Xue. "Overexpression of klotho suppresses growth and... <1%

31 Internet
journal.ugm.ac.id <1%

32 Internet
tessera.spandidos-publications.com <1%

33 Internet
watermark.silverchair.com <1%

34 Publication
Chrismis Novalinda Ginting, I Nyoman Ehrich Lister, Ermi Girsang, Wahyu Widow... <1%

35 Publication
Mitochondria as Targets for Phytochemicals in Cancer Prevention and Therapy, 2... <1%

36 Publication
Moshe Giladi, Mijal Munster, Rosa S. Schneiderman, Tali Voloshin et al. "Tumor tr... <1%

37 Publication
Xin Lu, Xisheng Weng, Zheng Li, Bo Yang, Jun Qian, Yue Huang. "Protective role of... <1%

38 Publication
J W Zimmerman. "Cancer cell proliferation is inhibited by specific modulation freq... <1%

39

Publication

Junjie Niu, Yibao Sun, Qiaoge Guo, Dongju Niu, Bo Liu. "Serum miR-95-3p is a diag... <1%

40

Publication

Marlina Marlina, Armenia Armenia, Rizki Rahmadian, Jenifer Kiem Aviani et al. "C... <1%

41

Publication

Yuan Fang, Xiao-yan Xu, Yubang Shen, Jiale Li. "miR-23a-3p and miR-23a-5p target... <1%

42

Publication

Rahaba Marima, Egnesious Sambo, Ntombikayise Xelwa, Olalekan Fadebi et al. "P... <1%

43

Internet

repository.maranatha.edu <1%

44

Internet

etheses.whiterose.ac.uk <1%

Non-contact Electric Field Exposure Provides Potential Cancer Therapy through p53-Independent Proliferation Arrest and Intrinsic Pathway Apoptosis Induction in MG-63 Cell Lines

Omat Rachmat Hasbullah^{1*}, Ilma Fiddiyanti², Dewi Ratih Handayani², Endang Sutedja³, Darmadji Ismono³, Nucki Nursjamsi Hidajat³, Wahyu Widowati⁴, Ervi Afifah⁵, Hanna Sari Widya Kusuma⁵, Rizal Rizal^{5,6}, Firman Alamsyah⁷, Warsito P. Taruno⁷

¹Faculty of Medicine, University of Sultan Ageng Tirtayasa, Serang, Banten 42124, Indonesia

²Faculty of Medicine, University of Jenderal Achmad Yani, Cimahi 40531, Indonesia

³Faculty of Medicine, Universitas Padjadjaran, Jatinangor 45363, Indonesia

⁴Faculty of Medicine, Maranatha Christian University, Bandung 40164, Indonesia

⁵Biomolecular and Biomedical Research Centre, Aretha Medika Utama, Bandung 40163, Indonesia

⁶Biomedical Engineering, Department of Electrical Engineering, Faculty of Engineering, Universitas Indonesia, Depok 16424, Indonesia

⁷Center for Medical Physics and Cancer Research Ctech Labs Edwar Technology, Tangerang 15143, Indonesia

ARTICLE INFO

Article history:

Received May 20, 2022

Received in revised form May 29, 2022

Accepted October 28, 2022

KEYWORDS:

Cancer,
p53,
p21,
MDM2,
Caspase 8,
Caspase 9

ABSTRACT

Osteosarcoma is a highly malignant primary tumor on bone that mainly attacks children and young adolescents. Until now, osteosarcoma therapy still combines some high costs and invasive therapy modalities that may give side effects, such as pain and nausea. Our previous studies suggested that non-contact electric field has anti-proliferative effect on breast cancer cells, *in vitro* and *in vivo*. In this study, we were interested studying alternating current electric field effects on osteosarcoma cells progression as well as its potential cytotoxic effects. MG-63 human osteosarcoma cells were cultured and treated with 200 kHz for 6 days. Several genes of interest including p53, p21, MDM2, caspase-3, caspase-8, and caspase-9 were analyzed using real-time qPCR method. Apoptotic index was measured using flow-cytometry assay. Apoptosis was observed through p53-independent p21 pathway ($p = 0.011$). Cells undergoing apoptosis through internal pathways were shown by the increase of caspase-3 ($p = 0.015$) and caspase-9 ($p = 0.001$) levels, but not caspase-8 ($p = 0.080$). This treatment has successfully reduced the number of living osteosarcoma cells by 14.7% ($p = 0.000$) and increased cell death up to 4.26% ($p = 0.055$). Apoptotic index was markedly increased to 16% ($p = 0.001$). 200 kHz non-contact electric field exposure can disrupt osteosarcoma progression through disruption of normal cell cycle via p53-independent p21 pathway and induction of apoptosis.

1. Introduction

Osteosarcoma is a highly malignant primary tumor on bone that mainly affects children and young adolescents (Solomon *et al.* 2010). In Indonesia, osteosarcoma was recorded to account for 22–25% of all bone and joint cancers (Hutagalung 2005; Ismono 2003). Osteosarcoma cells are the most significant inherent determinant for a bad prognosis in osteosarcoma patients because they are more prone

to spreading and developing metastases (Bielack *et al.* 2002). The majority of the time, osteosarcoma spreads to the lungs, while it can also happen to bone and lymph nodes, the progression of osteosarcoma is marked by aggressive growth and local recurrence, resulting in low life expectancy caused by metastasis to lungs in more than 50% cases. So far, osteosarcoma therapy still combines some therapy modalities, including radiation therapy, immunotherapy, chemotherapy which uses chemotherapeutic agent, biological agent or neoadjuvant therapy, and surgical treatment including limb salvage and amputation (Jaffe *et al.* 2013). However, these therapies are

* Corresponding Author

E-mail Address: dr_omatr@yahoo.com

high costs and can cause side effects, such as pain and nausea, and may also affect the healthy cells (Agarwal and Nayak 2012; Chen *et al.* 2013; Dicarpio *et al.* 2003; Hutagalung 2005; Harisson *et al.* 2018; Jaffe *et al.* 2013; Yang 2005).

Several alternative therapies have been proposed to treat osteosarcoma. Numerous cancer medicines that target cancer cells directly and specifically are under clinical development, e.g., multi-target drugs for receptor tyrosine kinase inhibitors, cell cycle and DNA repair proteins, or through indirect targeting of microenvironment modulation, such as macrophage regulator and loaded T lymphocytes (Brown *et al.* 2018). However, not only chemical drugs development, but several physical therapies have also been proposed, such as phototherapy (Yu *et al.* 2017), cold atmospheric plasma (Gumbel *et al.* 2017), ultrasound (Agnese *et al.* 2019), magnetic and water-based hyperthermia (Herea *et al.* 2018), and pulse electromagnetic field (Muramatsu *et al.* 2017). Having antimitotic effect on cells that ultimately results in cell death (Kirson *et al.* 2004; McCaig *et al.* 2005; Rachmat *et al.* 2017), currently, alternating current, low intensity, intermediate frequency electric fields, have been revealed as a novel cancer treatment (Hottinger *et al.* 2016; Kirson *et al.* 2004). In our previous study, low intensity (18 Vpp) and intermediate frequency (100 kHz) electric fields exposure has shown anti-proliferative effect on breast cancer cells, *in vitro* and *in vivo* (Alamsyah *et al.* 2015), and was successfully inhibited in the proliferation of HeLa and oral cancer cells (Mujib *et al.* 2017). This intermediate frequency is used to treat cancer because it specifically targets cancer cells without affecting normal cells. Furthermore, we develop non-contact electric fields to avoid dermatitis or other injuries due to direct contact between the electrodes and the skin of animal model, as reported by Kirson *et al.* (2007).

Apoptosis is the main barrier of oncogenesis and tumor suppressor protein p53 is the main key for the regulation of apoptosis and carcinogenesis which is mediated by two main pathways, extrinsic and intrinsic pathways. In extrinsic pathway, caspase 8 activation occurs to induce apoptosis, whereas in the intrinsic pathway, there is a role of mitochondrial protein in caspase 9 activation to induce apoptosis. Both mechanisms resulted in pro-caspase 3 cleavage to become active caspase 3, the main executioner in the apoptosis. Caspase-3 causes chromosomal DNA degradation and induces cytoskeletal reorganization

and cell body disintegration (Fiandalo and Kyprianou 2012; Lahiry *et al.* 2010; Maximov and Maximov 2008; Olsson and Zhivotovsky 2011; Parissh *et al.* 2013; Pirnia *et al.* 2002; Silke and Meier 2013). Protein p53 is an important factor that inhibits tumor progression by activating p21 protein that regulates cell cycle progression (Maximov and Maximov 2008). Mutations of p53 gene is the commonest mutation in more than 50% of cases of malignancy especially in osteosarcoma (Yao *et al.* 2014). The p53 protein in normal cells remains low so that cells keep growing and proliferating (Bai and Zhu 2006). Low p53 protein concentration is caused by the regulation of Murine Double Minute 2 (MDM2) oncoprotein. In most cases of osteosarcoma, MDM2 gene is overexpressed and this is related to a poor clinical prognosis (Hou *et al.* 2019), MDM2 functions as E3 ubiquitin ligase and is responsible for the continued degradation of p53 in normal cells.

Based on the backgrounds, intermediate frequency and low intensity electric field exposure can be a prospective, less invasive cancer therapy. As far as we know, this kind of novel treatment has not been done on osteosarcoma cancer cells. Based on the backgrounds, intermediate frequency and low intensity electric field exposure can be a prospective, less invasive cancer therapy. Thereby we were interested in studying the effect of this therapy on MG-63 cell line human osteosarcoma *in vitro* through the evaluation of apoptotic markers p53, p21, caspase 3, caspase 8, caspase 9, and MDM2 expression.

2. Materials and Methods

2.1. Cell Culture

MG-63 cell lines (ATCC®CRL-2522), provided by Aretha Medika Utama, Biomolecular and Biomedical Research Center, Bandung, West Java, Indonesia, were cultured in complete medium (Minimum Essential Medium- α (MEM- α) (Biowest L0475) + 10% Fetal Bovine Serum (FBS) (Biowest S181H) + 1% penicillin streptomycin (Biowest L0022) in 37 °C and 5% CO₂ conditions. After reaching 80 to 90% confluence, cells were harvested for treatments (Rachmat *et al.* 2017).

2.2. Treatments

Cells were harvested with 1 mL trypsin EDTA and centrifuged at 2,000 rpm for 5 minutes. Supernatant was discarded and pellet was suspended in 1 mL MEM- α medium. Cells were inserted into a 6-well

plate at 2×10^6 cells/well density. Complete medium was added as much as 3 ml and incubated for 24 hours. After 24 hours and the cells were already attached, the medium was replaced. The culture plates were positioned between two capacitive electrodes that were connected to an oscillator (Figure 1) that was provided by Centre for Medical Physics and Cancer Research Ctech Labs Edwar Technology (Alamsyah *et al.* 2015; Mujib *et al.* 2017). Later, the electrodes and plates were placed in incubator at 37°C and 5% CO₂. Cells were treated with 200 kHz for 6 days based on previous research with medium replacement every two days (Rachmat *et al.* 2017).

2.3. RNA Isolation and cDNA Synthesis

Aurum Total RNA Mini kit (Bio-Rad, 1708841) [California, USA] was used to perform RNA isolation. All procedures were referred to Afifah

et al. (2019) and Widowati *et al.* (2020a) protocol. The iScript cDNA synthesis kit (Bio-Rad, 1708890) [California, USA] was used on performing cDNA synthesis according to the manufacturer's protocol. Temperature reaction for cDNA synthesis was 25°C for 5 minutes; reverse transcription at 42°C for 30 minutes; RT Inactivation at 85°C for 5 minutes; and the last hold at 4°C for 5 min (Afifah *et al.* 2019; Widowati *et al.* 2020a).

2.4. Gene Expression Measurement with Real-Time PCR

The qPCR reaction was done using SsoFast Eva Green Supermix kit according to manufacturer's protocols (BioRad, 172-5200) [California, USA] which contained: 5 µl evagreen, 2 µl each reverse and forward primers (See Table 1), 2 µl Nuclease



Figure 1. Capacitive electrodes as electrical field generator. A microplate was flanked between two capacitive electrodes that were connected to low intensity oscillator to generate the alternate electric field

Table 1. Primer sequence and product size of casp9, casp8, casp3, MDM2, p21, and p53 genes

Primer	Forward	Reverse	Size (bp)
casp9	5'-CAT GCT CAG GAT GTA AGC CA-3'	5'-AGG TTC TCA GAC CGG AAA CA-3'	161
casp8	5'-AGG CCA GAT CTT CAC TGT CC-3'	5'-GGT CAC TTG AAC CTT GGG AA-3'	114
casp3	5'-AGA ACT GGA CTG TGG CAT TGA G -3'	5'-GCT TGT CGG CAT ACT GTT TCA G-3'	191
mdm2	5'-CTG ATC CAA CCA ATC ACC TG-3'	5'-AAG CCT GGC TCT GTG TGT AA-3'	130
p21	5'-CAG CGA CCT TCC TCA TCC A-3'	5'-GAC TCC TTG TTC CGC TGC TAA-3'	71
p53	5'- CCA AGG CGC ACT GGC CTC ATC -3'	5'-CAG AGG CTG GGG CAC AGC AGG -3'	205

free water, 1 μ l cDNA samples. The master mix was transferred to piko real plate and incubated in the qPCR machine (ThermoScience) [Massachusetts, USA] with the following stages: pre-denaturation at 95°C for 5 minutes, denaturation at 95°C for 1 minute, annealing at 50°C (for all primers) for 40 seconds, elongation at 72°C for 1 minute, post-elongation at 72°C for 5 minutes, melt curve 55–95°C, and hold at 4°C (Afifah *et al.* 2019; Widowati *et al.* 2020a). This stage was done in triplicate. Primer sequences and product sizes are shown in Table 1. Gene expression was measured as relative copy number compared to control using $\Delta\Delta CT$ method (Arya *et al.* 2005).

2.5. Apoptotic Indexes

Approximately 5×10^5 to 2×10^6 cells were taken and centrifuged at 1,600 rpm, at 4°C for 5 minutes. The supernatant was discarded, while the pellet was mixed with 1 mL of PBS (Gibco, 14200075) [Thermo Fisher Scientific, Massachusetts, USA] contain 5% FBS (Biowest, S181H) [Biowest, Nuaile, France]. Samples were then centrifuged at the same condition. Supernatant was removed and 1 ml binding buffer from Annexin V Fluos Kit (Sigma Aldrich, ab14085) [Sigma Aldrich, Missouri, USA] were added to the pellet. Then 5 ml of PI and FITC was added in the sample and it was incubated at 4°C for approximately 30–40 minutes (Gloushankova *et al.* 2017). Samples were later analyzed in MACSquant Flowcytometry (Miltenyi Biotech) [Bergisch Gladbach, Germany] (Lister *et al.* 2020; Widowati *et al.* 2020b).

2.6. Statistical Analysis

Independent t-test was used to compare significance between control and treatments group at confidence interval 95% ($p = 0.05$).

3. Results

3.1. Non-contact Electric Field Alters Cell Morphology

The alternating current electric field alters cells' morphology. The fibroblastic, spindle shaped

osteosarcoma had changed into spherical shaped cells. As also can be seen from (Figure 2), the gap between cells was seen at all treated groups.

3.2. Non-contact Electric Field May Causes Cell Cycle Arrest

The 200 kHz frequency of alternate current electric field in our experiment had caused significant changes on p21 gene and MDM2 gene expression (Table 2). The expression of p21 gene increased significantly compared to control which indicated increase inhibitory activity in cell cycle progression. Due to the MDM2 gene's role as a p53 negative regulator, its expression was lower than that of the control. Cell cycle arrest and apoptosis result from an increase in p53 levels, which also increases p53 transcriptional activity. Insignificant variations in p53 expression levels imply potential alternate p21 regulatory mechanisms protein activity that implicated increased p21 response and activity. However, insignificant changes in p53 expression level indicated that there might be an alternative pathway in p21 regulation.

3.3. Non-contact Electric Field Induces Apoptosis through Intrinsic Pathway

Electric field exposure had caused decrease in MG-63 human osteosarcoma cell survival by 14.7% ($p = 0.000$), referring to the percentage of live cells. The dead of the cells in treatment group was thought mainly due to apoptosis (Figure 3). The 200 kHz frequency of alternate current electric field had caused significant difference increase of apoptotic index from 1.95(1.77–2.07) to 17.53 (17.36–18.91) ($p = 0.001$), which means that the electric field induced cell apoptotic activity (Table 3). Caspase 9 expression difference significantly ($p = 0.001$), while changes in caspase 8 are insignificant ($p = 0.080$), indicates that the cells are mainly induced to undergo apoptotic activity through intrinsic pathway (Table 2). difference in caspase 9 expression implicated in difference of caspase 3 as the apoptotic executioner by four-fold ($p = 0.015$).




Figure 2. Electrical field reduces MG-63 alters cells' morphology. The figure above shows the MG-63 cell population through an inverted microscope at 6 days incubation at $20 \times 20,000$ magnification. The non-treated cells show triangular spindle shape morphology. The cells undergone massive proliferation and already reached 100% confluence at 6 days in the control group. Electrical field induction causes alteration on cells' morphology where they seem to shrink and have spherical morphology and show the gap between cells in the treatment group

Table 2. Effect of 200 kHz electrical field induction to p53, p21, MDM2, casp3, casp8, casp9 relative gene expression in MG-63 human osteosarcoma cells. Data was presented mean \pm standard deviation. The single star (*) show statistically significant differences based on the t-test ($p<0.05$)

Variable	Group		Significance (p-value)
	200 kHz treated	Control	
p53 Expression			
Mean (SD)	1.37 (0.06)	1 (0.0)	0.198
Median (range)	1.36 (1.31-1.43)	1 (1-1)	
p21 Expression			
Mean (SD)	9.96 (0.06)	1 (0.0)	0.011*
Median (range)	9.65 (11.71-8.51)	1 (1-1)	
MDM2 Expression			
Mean (SD)	0.67 (0.12)	1 (0.0)	0.065
Median (range)	0.63 (0.57-0.81)	1 (1-1)	




Figure 3. Percentage of MG-63 human osteosarcoma cell conditions with or without 200 kHz electrical induction treatment. The graph shows the proportion of live cells, dead cells, cells that underwent apoptosis, and necrotic cells based on the Flow cytometry result. Electrical induction can reduce MG-63 human osteosarcoma cells survival and induce them to undergo apoptosis resulted in cell death. Data was presented mean \pm standard deviation. The single star (*) show statistically significant differences based on the t-test ($p<0.05$)

Table 3. Effect of 200 kHz alternate current in MG-63 human osteosarcoma cells apoptotic activity. Data was presented mean \pm standard deviation. The single star (*) show statistically significant differences based on the t-test ($p<0.05$)

Variable	Group		Significance (p-value)
	200 kHz treated	Control	
Caspase 3 Expression			
Mean (SD)	4.05 (0.96)	1 (0.0)	0.015*
Median (range)	4.08 (3.07-4.99)	1 (1-1)	
Caspase 8 Expression			
Mean (SD)	1.55 (0.34)	1 (0.0)	0.080
Median (range)	1.47 (1.25-1.93)	1 (1-1)	
Caspase 9 Expression			
Mean (SD)	8.71 (1.69)	1 (0.0)	0.001*
Median (range)	8.82 (6.96-10.34)	1 (1-1)	
Apoptotic Index			
Mean (SD)	17.93 (0.85)	1.93 (0.15)	0.001*
Median (range)	17.53 (17.36-18.91)	1.95 (1.77-2.07)	

4. Discussion

200 kHz non-contact electric field is the effective frequency to treat osteosarcoma and the therapy should be performed for at least six days to give significant effect (Rachmat *et al.* 2017). As indicated in (Figure 2) the TTFields causes spindle shaped osteosarcoma cells to undergo morphological changes to sphere-like shapes with gap between cells during six days of treatment. The morphological changes indicate cell stress, which is caused by microtubule and actin filament network misalignments. These cytoskeletons also playing roles in maintaining cells adherent junctions and junction associated proteins such as cadherin/catenin that attach to F-actin which are maintained by microtubules (Gloushankova *et al.* 2017; Kasioulis *et al.* 2017), microtubule and actin filaments give mechanical strength and membrane stiffness to the cells (Kelliher *et al.* 2019).

Mitotic aberrations related to centrosome and mitotic spindle defects causes premature p53-p21 and p21-p16 premature senescence phenotype (Dikovskaya *et al.* 2015; Schmidt *et al.* 2010). Our result shows that p21 is significantly upregulated due to electrical field exposure, however p53 is not. This may indicate that there is a mechanism of p53-independent p21 upregulation pathway, although it has been known that p21 is p53 target. MELK (Maternal Embryonic Leucine Zipper Kinase) inhibition could cause cell cycle arrest at G1 stage through p21 expression in p53-knockdown cancer cells. Further investigation demonstrated two transcriptional regulators of p21, FOXO (Fork head box O) -1 and -3 which were de-activated through phosphorylation

by MELK, is upregulated in MELK-knock down cells (Matsuda *et al.* 2017). Another proposed mechanism is upregulation of p21/p27 through autophagy-mediated downregulation of F box protein Skp2 and p62/SQSTM1 expression independent of p53 status due to Quinacrine induction in ovarian cancer (Jung *et al.* 2018). Our data also shows that MDM2 is not significantly downregulated due to electrical field exposure. MDM2 is a protein that is responsible in Akt-MDM2-p53 signaling pathway in p21 regulation (Ignacio *et al.* 2018). The thing that drives apoptosis is from p53 signaling which transcribes and activates a large number of genes so that it undergoes apoptosis which in turn leads to tumorigenesis that causes DNA damage. a study conducted by Mujib *et al.* (2017) regarding the p53 expression of cancer cells under exposure to a low-intensity and medium-frequency non-contact electric field, found that 80-85% of positively stained cancer cells prevented cell division through p53 protein in oral cultured cells and HeLa. From (Figure 3) necrotic index is significantly higher ($p = 0.003$) in non-treated cells compared to treated cells. Induced by oxygen and glucose deprivation (OGD) and beginning with cell swelling, leading to membrane rupture and release of cytoplasmic contents into the extracellular space, necrosis has been considered as an accidental and genetically unprogrammed form of cell death. Cells in the inner regions of solid tumors (including osteosarcoma) display higher OGD because of insufficient blood supply. Necrosis has a promotion potential by releasing Damage-Associated Molecular Pattern molecules including HMGB1 (High Mobility Group Box1) which resulted in enhanced inflammatory responses, reactive oxygen formation, and metastasis

39 through the release of proinflammatory cytokines by activating NF κ B and inflammasomes pathway44. While our result shows that non-contact electric field exposure might potentially inhibit necrosis while promoting apoptosis, potentially, the necrosis inhibition in 200 kHz electric field treated cells is caused by inhibition of miR-23a-3p expression (Dong et al. 2017). The mechanism probably lies in miR-23a-3p because there is a sharing pathway between miR-23a-3p and pro-apoptotic protein bax (Grieco et al. 2017).

14 Our result suggests that non-contact electric field exposure causes intrinsic pathway apoptosis, marked by significant higher of casp3 and casp9 expression. Intrinsic pathway is a non-receptor-mediated pathway that cause intracellular cascades within the cell and is initiated within mitochondria. While some studies have suggested that electrical pulse stimulation may cause membrane permeabilization (Nuccitelli et al. 2007; Silve et al. 2015, 2016), the inner mitochondrial membrane could change because of pore opening of mitochondrial permeability transition, mitochondrial transmembrane potential loss, and pro-apoptotic proteins release from the intermembrane space into the cytosol (Elmore 2007). We hypothesized that the same mechanism might cause mitochondrial membrane permeabilization that leads to initiation of apoptosis intrinsic pathway.

31 Control and regulation of the mitochondrial-mediated apoptotic event are initiated by members of the Bcl-2 protein family. Bax (Bcl2-associated X protein), one of the family members, plays a role in cytochrome c release that leads to activation of pro-caspase-9. Later caspase-9 will cleave caspase-3 as the executioner. This protein will act through activation of CAD (Caspase-Activated DNase) in chromosomal DNA degradation. In addition to the tumor suppressor protein p53 having a critical role in the Bcl-2 proteins family regulation (Elmore 2007), Caspase 3 induces cytoskeletal reorganization and the cell disintegration into apoptotic bodies. Tumor treating fields (TTFields) prevent DNA damage repair by hindering homologous recombination repair, which may enhance Bax expression and enhance intrinsic pathway apoptosis (Giladi et al. 2017). The insignificance between p53 level in electric field treated tumor cells and control may suggest that there might be another mechanism of intrinsic pathway apoptosis which is p53-independent.

1 1 In osteosarcoma patients, the microRNA-95-3p is upregulated. When compared to normal controls, miR-95-3p expression was decreased in both high-grade and low-grade osteosarcoma tissues (Novello et al. 2013). Reduced expression of miR-95-3p is a statistically significant risk factor that lowers overall survival in patients with osteosarcoma and works as an anti-oncogenic miRNA in osteosarcoma (Niu et al. 2016). The miRNA-95-3p expression is probably inhibited during electric field exposure, in osteosarcoma cells, micro-RNA-95-3p inhibition increases caspase-3 and caspase-9 activities and Bax/Bcl-2 protein expression, and eventually suppresses cell growth and induces apoptosis. MiRNA-95-3p suppression also gives anticancer effects on cell growth by suppressing TGF- β and p-Smad2 while inducing p21 protein expression and suppressing cyclin D1 protein (Zhao et al. 2018).

32 33 9 35 From our result and literature review, we proposed the possible mechanism of TTField induction in inhibiting osteosarcoma progression as shown in (Figure 4). The alternating current can affect osteosarcoma cells through several mechanisms: 1) It causes mitotic aberration that causes p21 senescence phenotype through p53-independent pathway. This pathway is probably initiated by MELK inhibition that causes FOXO1 and FOXO3 transcription factor activation which upregulates p21. The upregulation of p21 resulted in inhibition of cyclin D1 that causes G1/S phase progression delay (senescence); 2) miR-23a-3p inhibition that resulted in necrosis inhibition and apoptosis progression; 3) inhibition of miR-95-3p that upregulates p21 and Bax, caspase9, and caspase3, marker proteins of apoptosis intrinsic pathway; 4) increased mitochondrial permeability that resulted in intrinsic pathway upregulation. In the intrinsic pathway apoptosis, protein Bax will cause cytochrome c release and together with caspase9 will create apoptosome that will result in caspase3 activation. Caspase3 later will cause cytoskeletal disorganization and DNA degradation through CAD activation; and 5) Inhibition of homologous repair that causes DNA repair delay.

21 17 36 2 In conclusion, non-contact electric field exposure can disrupt osteosarcoma progression through disruption of cell morphology and adhesion between cells. Osteosarcoma treated with 200 kHz for 6 days gave the cell senescence through p53-independent p21 pathway. Cells undergone apoptosis through




Figure 4. The proposed mechanism on how alternating current can induce cell-cycle arrest and apoptosis intrinsic pathway in osteosarcoma cells

intrinsic pathway by the marked increase of casp-3 and casp-9 levels. This treatment had successfully reduced the number of living osteosarcoma cells by 14.7% and increased cell death by 4.26%. The non-contact electric field is an attractive candidate to serve as a novel, non-invasive treatment for osteosarcoma.

Conflict of Interest

The authors declare that they have no competing interests.

Acknowledgments

We extend our gratitude to Agung Novianto, Viranda andria Yuninda, Anis Syahbani Muthmainnah and Afif Yati from Aretha Medika Utama, Biomolecular and Biomedical Research Center, Bandung, West Java, Indonesia for their technical support.

References

Afifah, E., Mozef, T., Sandra, F., Arumwardana, S., Rihibih, D.D., Nufus, H., Rizal, R., Amalia, A., Bachtiar, I., Murti, H., Widowati, W., 2019. Induction of matrix metalloproteinases in chondrocytes by interleukin IL-1 β as an osteoarthritis model. *J. Math. Fundam Sci.* 5, 1–9. <https://doi.org/10.5614/j.math.fund.sci.2019.51.2.1>

Agarwal, M.G., Nayak, P., 2012. Limb salvage for osteosarcoma: current status with a review of literature In: Agarwal M (Eds.). *Osteosarcoma*. Shanghai: InTech. pp. 109–136.

Agnese, V., Costa, V., Scoarughi, G.L., Corso, C., Carina, V., De Luca, A., Bellavia, D., Raimondi, L., Pagani, S., Midiri, M., Stassi, G., 2019. Focused ultrasound effects on osteosarcoma cell lines. *BioMed Res. Int.* 2019, 1–14. <https://doi.org/10.1155/2019/6082304>

Alamsyah, F., Ajrina, I.N., Dewi, F.N.A., Iskandriati, D., Prabandari, S.A., Taruno, W.P., 2015. Antiproliferative effect on electric fields on breast tumor cells *in vitro* and *in vivo*. *Indones. J. Cancer Chemoprevention*. 6, 71–77. <https://doi.org/10.14499/indonesianjcanchemoprev6iss3pp71-77>

Arya, M., Shergill, I.S., Williamson, M., Gommersall, L., Arya, N., Patel, H.R., 2005. Basic principles of real-time quantitative PCR. *Expert Rev. Mol. Diagn.* 5, 209–219. <https://doi.org/10.1586/14737159.5.2.209>

Bai, L., Zhu, W.G., 2006. p53: Structure, function and therapeutic applications. *J. Cancer Mol.* 2, 141–153.

Bielack, S.S., Kempf-Bielack, B., Delling, G., Exner, G.U., Flege, S., Helmke, K., Kotz, R., Salzer-Kuntschik, M., Werner, M., Winkelmann, W., Zoubek, A., 2002. Prognostic factors in high-grade osteosarcoma of the extremities or trunk: an analysis of 1,702 patients treated on neoadjuvant cooperative osteosarcoma study group protocols. *J. Clin. Oncol.* 20, 776–790. <https://doi.org/10.1200/JCO.2002.20.3.776>

Brown, H.K., Schiavone, K., Gouin, F., Heymann, M.F., Heymann, D., 2018. Biology of bone sarcomas and new therapeutic developments. *Calcif. Tissue Int.* 102, 174–195. <https://doi.org/10.1007/s00223-017-0372-2>

Chen, Y., Yang, Y., Wang, C., Shi, Y., 2013. Radiographic response to neoadjuvant therapy and its impact on scope of surgery and prognosis in stage IIIB/III soft tissue sarcomas. *BMC Cancer.* 13, 1–9. <https://doi.org/10.1186/1471-2407-13-591>

Dikovskaya, D., Cole, J.J., Mason, S.M., Nixon, C., Karim, S.A., McGarry, L., Clark, W., Hewitt, R.N., Sammons, M.A., Zhu, J., Athineos, D., 2015. Mitotic stress is an integral part of the oncogene-induced senescence program that promotes multinucleation and cell cycle arrest. *Cell Rep.* 12, 1483–1496. <https://doi.org/10.1016/j.celrep.2015.07.055>

Dong, Y., Li, T., Li, Y., Ren, S., Fan, J., Weng, X., 2017. MicroRNA-23a-3p inhibitor decreases osteonecrosis incidence in a rat model. *Mol. Med. Rep.* 16, 9331–9336. <https://doi.org/10.3892/mmr.2017.7808>

Elmore, S., 2007. Apoptosis: a review of programmed cell death. *Toxicol. Pathol.* 35, 495–516. <https://doi.org/10.1080/01926230701320337>

Fiandalo, M.V., Kyriyanou, N., 2012. Caspase control: protagonists of cancer cell apoptosis. *Exp. Oncol.* 34, 165–175.

Giladi, M., Munster, M., Schneiderman, R.S., Voloshin, T., Porat, Y., Blat, R., Zielinska-Chomej, K., Hågg, P., Bomzon, Z.E., Kirson, E.D., Weinberg, U., 2017. Tumour treating fields (TTFields) delay DNA damage repair following radiation treatment of glioma cells. *Radiat. Oncol.* 12, 1–13. <https://doi.org/10.1186/s13014-017-0941-6>

Gloushankova, N.A., Rubtsova, S.N., Zhitnyak, I.Y., 2017. Cadherin-mediated cell-cell interactions in normal and cancer cells. *Tissue Barriers.* 5, 1–15. <https://doi.org/10.1080/21688370.2017.1356900>

Grieco, F.A., Sebastiani, G., Juan-Mateu, J., Villate, O., Marroqui, L., Ladrière, L., Tugay, K., Regazzi, R., Bugiani, M., Marchetti, P., Dotta, F., 2017. MicroRNAs miR-23a-3p, miR-23b-3p, miR-149-5p regulate the expression of proapoptotic BH3-only proteins DP5 and PUMA in human pancreatic b-cells. *Diabetes.* 66, 100–112. <https://doi.org/10.2337/db16-0592>

Gumbel, D., Bekeschus, S., Gelbrich, N., Napp, M., Ekkernkamp, A., Kramer, A., Stope, M.B., 2017. Cold atmospheric plasma in the treatment of osteosarcoma. *Int. J. Mol. Sci.* 18, 1–13. <https://doi.org/10.3390/ijms18092004>

Harrison, D.J., Geller, D.S., Gill, J.D., Lewis, V.O., Gorlick, R., 2018. Current and future therapeutic approaches for osteosarcoma. *Expert Rev. Anticancer Ther.* 18, 39–50. <https://doi.org/10.1080/14737140.2018.1413939>

Hutagalung, E.U., 2005. Overview of bone tumours in Cipto Mangunkusumo Hospital Jakarta management and problems. In: *Proceedings of 10th National Conference Pathology and Anatomy Professionals Associations in Indonesia.* Surabaya: National Conference Pathology and Anatomy Professionals Associations in Indonesia, pp. 24–51.

Herea, D.D., Danceanu, C., Radu, E., Labusca, L., Lupu, N., Chiriac, H., 2018. Comparative effects of magnetic and water-based hyperthermia treatments on human osteosarcoma cells. *Int. J. Nanomed.* 13, 5743–5751. <https://doi.org/10.2147/IJN.S174853>

Hottinger, A.F., Pachero, P., Stupp, R., 2016. Tumour treating fields: a novel treatment modality and its use in brain tumours. *Neuro-Oncol.* 18, 1338–1349. <https://doi.org/10.1093/neuonc/now182>

Hou, H., Sun, D., Zhang, X., 2019. The role of MDM2 amplification and overexpression in therapeutic resistance of malignant tumours. *Cancer Cell Int.* 19, 216–223. <https://doi.org/10.1186/s12935-019-0937-4>

Ignacio, R.M., Dong, Y.L., Kabir, S.M., Choi, H., Lee, E.S., Wilson, A.J., Beeghly-Fadiel, A., Whalen, M.M., Son, D.S., 2018. CXCR2 is a negative regulator of p21 in p53-dependent and independent manner via Akt-mediated MDM2 in ovarian cancer. *Oncotarget.* 9, 9751–9765. <https://doi.org/10.18632/oncotarget.24231>

Ismono, D., 2003. The profile of musculoskeletal tumour in Hasan Sadikin Hospital. In: *Proceedings Scientific Meeting and Workshop of Musculoskeletal Pathology.* Surabaya: Scientific Meeting and Workshop of Musculoskeletal Pathology, pp. 181–192.

Jaffe, N., Puri, A., Gelderblom, H., 2013. Osteosarcoma: evolution of treatment paradigms. *Sarcoma.* 2013, 1–7. <https://doi.org/10.1155/2013/203531>

Jung, D., Khurana, A., Roy, D., Kalogera, E., Bakkum-Gamez, J., Chien, J., Shridhar, V., 2018. Quinacrine upregulates p21/p27 independent of p53 through autophagy-mediated downregulation of p62-Skp2 axis in ovarian cancer. *Sci. Rep.* 8, 1–12. <https://doi.org/10.1038/s41598-018-20531-w>

Kasioulis, I., Das, R.M., Storey, K.G., 2017. Inter-dependent apical microtubule and actin dynamics orchestrate centrosome retention and neuronal delamination. *eLife.* 6, 1–31. <https://doi.org/10.7554/eLife.26215>

Kelliher, M.T., Saunders, H.A., Wildonger, J., 2019. Microtubule control of functional architecture in neurons. *Curr. Opin. Neurobiol.* 57, 39–45. <https://doi.org/10.1016/j.conb.2019.01.003>

Kirson, E.D., Gurvich, Z., Schneiderman, R., Dekel, E., Itzhaki, A., Wasserman, Y., Schatzberger, R., Palti, Y., 2004. Disruption of cancer cell replication by alternating electric fields. *Cancer Res.* 64, 3288–3295. <https://doi.org/10.1158/0008-5472.CAN-00-0083>

Kirson, E.D., Dbaly, V., Tovaryš, F., Vymazal, J., Soustiel, J.F., Itzhaki, A., Mordechovich, D., Steinberg-Shapira, S., Gurvich, Z., Schneiderman, R., Wasserman, Y., 2007. Alternating electric fields arrest cell proliferation in animal tumor models and human brain tumors. *Proc. Natl. Acad. Sci. U. S. A.* 104, 10152–10157. <https://doi.org/10.1073/pnas.0702916104>

Lahiry, L., Saha, B., Chakraborty, J., Adhikary, A., Mohanty, S., Hossain, D.M.S., Banerjee, S., Das, K., Sa, G., Das, T., 2010. Theaflavins target Fas/caspase-8 and Akt/pBad pathways to induce apoptosis in p53-mutated human breast cancer cells. *Carcinogen.* 31, 259–268. <https://doi.org/10.1093/carcin/bgp240>

Lister, I.N.E., Ginting, C.N., Girsang, E., Nataya, E.D., Azizah, A.M., Widowati, W., 2020. Hepatoprotective properties of red betel (*Piper crocatum* Ruiz and Pav) leaves extract towards H₂O₂-induced HepG2 cells via anti-inflammatory, antinecrotic, antioxidant potency. *Saudi Pharmaceutical Journal.* 28, 1182–1189. <https://doi.org/10.1016/j.jps.2020.08.007>

Matsuda, T., Kato, T., Kiyotani, K., Tarhan, Y.E., Saloura, V., Chung, S., Ueda, K., Nakamura, Y., Park, J.H., 2017. P53 independent p21 induction by MELK inhibition. *Oncotarget.* 8, 57938–57947. <https://doi.org/10.18632/oncotarget.18488>

Maximov, G.K., Maximov, K.G., 2008. The role of p53 tumour-suppressor protein in apoptosis and cancerogenesis. *Biotechnol. Biotechnol. Equip.* 22, 664–668. <https://doi.org/10.1080/13102818.2008.10817532>

McCaig, C.D., Rajnicek, A.M., Song, B., Zhao, M., 2005. Controlling cell behavior electrically: current views and future potential. *Physiol. Rev.* 85, 943–978. <https://doi.org/10.1152/physrev.00020.2004>

Mujib, S.A., Alamsyah, F., Taruno, W.P., 2017. Cell death and induced p53 expression in oral cancer, HeLa, and bone marrow mesenchyme cells under the exposure to noncontact electric fields. *Integr. Med. Int.* 4, 161–170. <https://doi.org/10.1159/000485186>

Muramatsu, Y., Matsui, T., Deie, M., Sato, K., 2017. Pulsed electromagnetic field stimulation promotes anti-cell proliferative activity in doxorubicine-treated mouse osteosarcoma cells. *In vivo.* 31, 61–68. <https://doi.org/10.21873/invivo.11026>

Niu, J., Sun, Y., Guo, Q., Niu, D., Liu, B., 2016. Serum miR-95-3p is a diagnostic and prognostic marker for osteosarcoma. *Springerplus.* 5, 1947. <https://doi.org/10.1186/s40064-016-3640-0>

Novello, C., Pazzaglia, L., Cingolani, C., Conti, A., Quattrini, I., Manara, M.C., Tognon, M., Picci, P., Benassi, M.S., 2013. miRNA expression profile in human osteosarcoma: role of miR-1 and miR-133b in proliferation and cell cycle control. *Int J Oncol.* 42:667–675. <https://doi.org/10.3892/ijo.2012.1717>

Nuccitelli, R., McDaniel, A., Anand, S., Cha, J., Mallon, Z., Berridge, J.C., Uecker, D., 2017. Nano-pulse stimulation is a physical modality that can trigger immunogenic tumour cell death. *J. Immunother. Cancer.* 5, 1–13. <https://doi.org/10.1186/s40425-017-0234-5>

Olsson, M., Zhivotovsky, B., 2011. Caspases and cancer. *Cell Death Differ.* 18, 1441–1449. <https://doi.org/10.1038/cdd.2011.30>

Parrish, A.B., Freel, C.D., Kornbluth, S., 2013. Cellular mechanisms controlling caspase activation and function. *Cold Spring Harb. Perspect. Biol.* 5, 1–24. <https://doi.org/10.1101/cshperspect.a008672>

Pirnia, F., Schneider, E., Betticher, D.C., Borner, M.M., 2002. Mitomycin C induces apoptosis and caspase-8 and -9 processing through a caspase-3 and Fas-independent pathway. *Cell Death Differ.* 9, 905–914. <https://doi.org/10.1038/sj.cdd.4401062>

Rachmat, O., Fidhiyanti, I., Sutedja, E., Ismono, D., Hidajat, N.N., 2017. Effect of electric fields on osteosarcoma (MG-63) cells. *Int. J. Sci. Res. Educ.* 5, 6513–6517.

Schmidt, S., Essmann, F., Cirstea, I.C., Kuck, F., Thakur, H.C., Singh, M., Kletke, A., Jänicke, R.U., Wiek, C., Hanenberg, H., Ahmadian, M.R., 2010. The centrosome and mitotic spindle apparatus in cancer and senescence. *Cell Cycle.* 9, 4469–4473. <https://doi.org/10.4161/cc.9.22.13684>

Silke, J., Meier, P., 2013. Inhibitor of apoptosis (IAP) proteins-modulators of cell death and inflammation. *Cold Spring Harb. Perspect. Biol.* 5, 1–19. <https://doi.org/10.1101/cshperspect.a008730>

Silve, A., Leray, I., Leguebe, M., Poignard, C., Mir, L.M., 2015. Cell membrane permeabilization by 12-ns electric pulses: Not a purely dielectric, but a charge-dependent phenomenon. *Bioelectrochemistry.* 106, 369–378. <https://doi.org/10.1016/j.bioelechem.2015.06.002>

Silve, A., Leray, I., Poignard, C., Mir, L.M., 2016. Impact of external medium conductivity on cell membrane electropermeabilization by microsecond and nanosecond electric pulses. *Sci Rep.* 6, 1–15. <https://doi.org/10.1038/srep19957>

Solomon, L., Warwick, D., Nayagam, S., 2010. *Apley's System of Orthopaedics and Fractures*, ninth ed. Hodder Arnold, London.

Widowati, W., Jasaputra, D.K., Wargasetia, T.L., Azizah, A.M., Subangkit, M., Lister, I.N.E., Ginting, C.N., Girsang, E., Faried, A., 2020a. Apoptotic potential of secretome from interleukin-induced natural killer cells toward breast cancer cell line by transwell assay. *HAYATI J. Biosci.* 27, 186–196. <https://doi.org/10.4308/hjb.27.3.186>

Widowati, W., Jasaputra, D.K., Sumitro, S.B., Widodo, M.A., Mozef, T., Rizal, R., Kusuma, H.S.W., Laksmitawati, D.R., Murti, H., Bachtiar, I., Faried, A., 2020b. Effect of interleukins (IL-2, IL-15, IL-18) on receptors activation and cytotoxic activity of natural killer cells in breast cancer cell. *African Health Sci.* 20, 822–832. <https://doi.org/10.4314/ahs.v20i2.36>

Yang, R.S., 2005. Limb salvage operations for patients with malignant bone tumours in the extremities. *Tzu Chi Med. J.* 17, 389–396.

Yao, D., Cai, G.H., Chen, J., Ling, R., Wu, S.X., Li, Y.P., 2014. Prognostic value of p53 alterations in human osteosarcoma: a meta-analysis. *Int. J. Clin. Exp. Pathol.* 7, 6725–6733.

Yu, W., Zhu, J., Wang, Y., Wang, J., Fang, W., Xia, K., Shao, J., Wu, M., Liu, B., Liang, C., Ye, C., 2017. A review and outlook in the treatment of osteosarcoma and other deep tumors with photodynamic therapy: from basic to deep. *Oncotarget.* 8, 39833–39848. <https://doi.org/10.18632/oncotarget.16243>

Zhao, X., Yang, Y., Xu, J., Luo, Y., Xin, Y., Wang, Y., 2018. Downregulation of microRNA-95-3p suppresses cell growth of osteosarcoma via CDKN1A/p21 expression. *Oncol. Rep.* 39, 289–297. <https://doi.org/10.3892/or.2017.6065>