



## *Canna indica* L and *Spirulina platensis* for Food Security

Nita Noriko<sup>1\*</sup>, Risa Swandari Wijihastuti<sup>1</sup>, Alina Primasari<sup>1</sup>, Analekta Tiara Perdana<sup>1</sup>, Hidayat Yorianta Sasaerila<sup>1</sup>

<sup>1</sup>Department of Biology, Faculty of Science and Technology, Universitas Al Azhar Indonesia (UAI), South Jakarta, Indonesia

Received: October 30, 2023

Revised: November 19, 2023

Accepted: December 25, 2023

Published: December 31, 2023

Corresponding Author:

Nita Noriko

[nitanoriko1707@gmail.com](mailto:nitanoriko1707@gmail.com)

DOI: [10.29303/jppipa.v9iSpecialIssue.5888](https://doi.org/10.29303/jppipa.v9iSpecialIssue.5888)

© 2023 The Authors. This open access article is distributed under a (CC-BY License)



**Abstract:** Climate change threatens to food security, because it decreases agricultural productivity. *C. indica* L can adapt to environment and produce white and red rhizomes. Dominant content both of rhizomes are carbohydrates which increase protein as nutrition by adding *S. platensis*. The research aims to determine potential of composite flour of *C. indica* L and *S. platensis* as alternative healthy foodstuff for food security. The method consists of formulation composite flour, proximate tests, determining amino and fatty acids, in vitro growth tests of *Lactobacillus*, cookies and meatball formulations. The result showed composite flour from red rhizome (RRCS) contains 11% water, 21.745% protein, 0.5% lipid, 9.99% dietary fiber, 57.81% carbohydrates, and 8.94% ash. While white rhizome (WRCS) contains, 8.75% water, 17.41 protein, 0.29% lipid, 9.54% dietary fiber, 67.601% carbohydrates and 5.94% ash. Amino and fatty acid appears in both composite flours were similar, except proline and oleic acid higher in RRCS. Lignoceric founded only in WRCS. In vitro test showed *Lactobacillus* (BAL) grew on modified MRSA from RRCS and WRCS. Both of composite flour processed into marketable cookies and meatballs formulations. Based on the result, *C. indica* L and *S. platensis* composite flour can be used as alternative food for food security.

**Keywords:** Change; Climate; Healthy Food; Security

### Introduction

Food security is defined in Law No. 18 of 2012 as the food availability for the state to persons in sufficient numbers and of good quality, that is safe and nutritious, but that should not violate religious rules, customs, and culture (Muliarta et al., 2023). Climate change is one of the factors threatening food security. The impact of climate change included drought, pest and disease resilience crops (Bvenura & Kambizi, 2022). Those conditions are the cause of agricultural productivity deduction. The simulation results of the annual crop productivity model due to climate change occurs in the middle of the 21st century. It shows a deduction in food security productivity globally approximately 3-12%. By the end of the 21st century will reach 11-25% (Wing et al., 2021). One of these phenomena already occur in West Nusa Tenggara Indonesia by lack of rice production (Khairulbahri, 2021). Other result shows that harvest corn production in China at time duration between 1979

to 2016 less of corn harvest or production by 1.7%. As facts happened the corn production became 5.19 kg 667 m<sup>-2</sup> every temperature increase 1° C (WU et al., 2021) the increase in temperature due to climate change that occurred in 1998 to 2017 also decreased corn production in Malang in 2018.

Lack of food production can be a cause of hunger. The World Food program explains that in 2020 there were 768 million people experiencing hunger (Syakirotin et al., 2022). Data shows that 95% of hunger that occurs is caused by lack of inadequate energy (Amelia et al., 2023). The main source of energy comes from plants that produce carbohydrates. This phenomenon shows indications of food insecurity that can affect nutritional status (Firna et al., 2018). Based on existing conditions, we need to find food source alternative which are can adapt with current climate change. This is in line with one of the government's steps in maintaining food security by diversifying food (Indrayani et al., 2023). In the growth of children,

### How to Cite:

**Example:** Susilawati, S., Doyan, A., Muliyadi, L., & Hakim, S. (2019). Growth of tin oxide thin film by aluminum and fluorine doping using spin coating Sol-Gel techniques. *Jurnal Penelitian Pendidikan IPA*, 1(1), 1-4. <https://doi.org/10.29303/jppipa.v1i1.264>

diversity of food consumption is needed to avoid stunting (Suhaimi et al., 2023). One alternative food source is jackfruit seed and oyster mushroom (Nirmalawaty et al., 2023). Other alternative food sources that are easy to grow and able to adapt to environment also need to be studied.

*C.indica* L. is a plant can adapt well to environmental such as intensity of sunlight and recommended for agroforestry practice (Sasaerila et al., 2021). *C. indica* L is native to South America and has spread throughout the world. This plant is widely cultivated in the tropics because it produces rhizome which is used as alternative food (Riandini et al., 2023)

The results of a study showed *C. indica* L contain 28.5% starch, 5.75% lipid, 4.72% crude protein, 25.1% fiber and vitamin with total estimated caloric values of 423/100 g dry weight (Ayusman et al., 2020a). *C. indica* L has two colors variation of the resulting rhizome, named as red and white. *C. indica* L starch belongs to the international standard quality range and can be used as confectioneries, noodles, thickening soups and sauces (Algar et al., 2019).

Based on nutrition content *C. indica* L has potential to use as alternative foodstuff. To increase protein levels on rhizome, *C. indica* L is proven by adding *S. platensis* which is a microalgae. This species contains chlorophyll a and phycobiliprotein that can live in saline conditions (Widawati et al., 2022). *S. platensis* can live on freshwater, in tropical weather and arid regions (Jung et al., 2019a) and able to adapt to the extreme environments such as pH and saline (Ismail et al., 2018).

Dry of *S. platensis* content 55-70% protein, 15-25% carbohydrate, 6-9% lipid, 8-10% dietary fiber, and 3-7% humidity (Jung et al., 2019b). *S. platensis* also contains mineral, vitamin, fiber, phycocyanin, pigments, unsaturated fatty acids, beta carotene, thiamine, nicotinamide, pyridoxine, riboflavin, fatty acid, phenolic, phycobilin protein, catechin hydrate, epicatechin, enzyme Superoxide Dismutase (SOD). Antibacterial, anticancer, antidiabetic, antioxidant, antihyperglycemic, antiinflammation dan antihypertension (Kamaludin & Holik, 2022) .

In addition to nutritional content, it is necessary to know the potential source of these foodstuffs as prebiotics. The importance of prebiotic in food as for health promotor supporting fermentation of intestinal microbiota. Fermentation will produce of Short Chain Fatty Acid (SCFA) that provides benefits for lipid, glucose, and immune homeostasis in conditions adequate intestinal microbiota (Silva et al., 2020). Existence of SCFA in intestine reduce the risk of colorectal cancer and inflammation in intestine (Campos-Perez & Martinez-Lopez, 2021). *S. platensis* from the Java Sea contains omega 6, gamma linoleic acid, carotenoid dan vitamin (Priyanti et al., 2022)

The obstacles to achieve food security is unacknowledgeable of health benefits alternative food sources in the community including process methods as various menus for daily consuming. However, composition of flour from *C. indica* L and *S. platensis* are not popular. As people do not know the benefits of consuming and processing into various menus which already common. Cookies and meatballs are favorable menus by market. The manufacturing processes from composite flour form *C. indica* L flour and *S. platensis* are unacknowledgeable. However, the biscuits manufacturing from wheat and Spirulina 2, 4, 6, and 5% was success (Saharan & Jood., 2017).

The aims of research are to determine potential of composite flour of *C. indica* L and *S. platensis* as alternative healthy foodstuff for food security in the face of climate change. The potency is based on nutritional considerations including the growth response of Lactobacillus in vitro as well as formulations for making cookies and meatballs

## Method

### Chemicals and instrumentation

Samples were prepared at chemistry laboratory of Al Azhar University, South Jakarta, Indonesia) and kept at room temperature. Afterwards proximate analysis was conducted. Fatty acids and amino acids composition was determined in the chemistry laboratory of the Center for Post-Harvest Research and Development, Bogor. In vitro growth of *L. casei* Shirota and *L. rhamnosus* was examined in the microbiology laboratory of Indonesia Al Azhar University.

### Steps of the research

The stages of research consist of 5 which are carried out sequentially (Figure 1).




Figure 1. The stages of research

### *C. indica* L Flour Preparation

Good-quality white and red *C. indica* L rhizomes were selected, peeled, washed, and sliced. These slices were immersed in sodium metabisulfite solution acted as an anti-browning agent. Then drained and sun dried for 3 days. The dried rhizomes were ground and sifted becoming flour.

### *S. platensis* Flour Preparation

*S. platensis* with Optical Density (OD) 0.5-0.7 was sub-cultured in a medium containing Nitrogen (N), Phosphate (P), and Potassium (K) (16:16:16) up to 0.75 g/L, and 1% NaOH added to maintain the pH at 9.5 for 12 days. The samples reached OD 0.7-1 harvested for further processing into flour. The flour prepared through flocculation, filtering, drying, and milling. A bio flocculant solution containing a mixture of 20 mL of vinegar + 0.2 g of chitosan + 180 mL of distilled water added to *S. platensis* in a ratio of 1: 100 (v/v). Afterwards, the flour filtered by filter paper. The left residue dried in an oven at of 45-70°C for ± 1 day. Dried sample grounded and spirulina flour was obtained.

### Composite Flour Preparation

*C. indica* L flour from red and white rhizomes and *S. platensis* flour mixed at a ratio of 10:1. This mixture stirred until homogeneous and flour was obtained.

### Composite Flour Proximate Composition

Composite Flour proximate composition was analyzed according to the Association of Official Methods of Analytical Chemists (AOAC, 2010) approved methods, including water, protein, lipid, ash, carbohydrate, and dietary fiber content.

### Determination of Fatty Acids Composition

A total of 2 mL of the lipid-extracted sample put into a test tube and 4.5 mL of 0.5 N NaOH added in methanol. It stirred, heated for 5 minutes at 50°C, then cool it down. Afterwards, 3 ml of BF<sub>3</sub> poured into the methanol and mixed using a vortex at 50°C for 5 minutes. Added Hexane and mixed it with vortex at 50°C for 5 minutes. Top layer of sample poured into a vial and injected into the GCMS machine.

### Determination of Amino Acids Composition

A sample of 0.05 g put into an Erlenmeyer flask mixed with 5 mL of 6 N HCl. It was hydrolysed and put in an oven at 115°C for 12 hours, filtered with filter paper. A 0.5 mL of it was transferred into a test tube. The sample dried using nitrogen gas mixed with 3 ml of 0.02 N HCl. Afterward, filtration carried out with Millipore 0.45 PTEE. A total of 20 µL sample extract poured into a vial tube followed by of 140 µL buffer isolate and 40 L reinforcing reagent. The sample put into the oven and injected into the HPLC for 45 minutes.

### In vitro growth of *L. casei* Shirota and *L. rhamnosus*

The MRSA medium used for vitro assays to grow *L. casei* Shirota and *L. rhamnosus*. Both composite flours applied as an additive to modify the medium. Glucose and peptone in the De Mann Rogosa and Shape Agar (MRSA) as substitute mixed with 20 g of *C. indica* L and

10 g of *S. platensis*. Based on the formulation, there were combinations of modified bacterial growth media.

*L. casei* Shirota grown in unmodified MRSA (UMW). Modified MRSA with the substitution of white rhizome flour (M-GCW), modified MRSA with the addition of *S. platensis* flour (M-PSW), as well as modified MRSA with the addition of white rhizome flour and *S. platensis* flour (M-GCW-PS).

*L. rhamnosus* grown in unmodified MRSA (UMR). Modified MRSA with substitution of red rhizome flour (M-GCR) then modified MRSA with the addition of *S. platensis* flour (M-PSR), as well as modified MRSA with substitution of *C. indica* L red rhizome flour and *S. platensis* flour (M-GCR-PS).

The growth medium set at a pH ranging from 6.4 to 6.6, while all media sterilized by autoclave and placed in 10 mL petri dishes. Stock cultures of *L. casei* Shirota and *L. rhamnosus* grown on the MRSA medium, then put into 10 mL of 0.85% NaCl and diluted. Both bacteria inoculated by spread plate method into each test medium with different formulations, and then incubated at 37°C aerobically for 48 hours. The total population in all media formulated as follows (Chouhan, 2015; Wulandari, 2023).

$$\text{Total population } \left(\frac{\text{CFU}}{\text{mL}}\right) = \text{Number of colonies} \times \frac{1}{\text{Dilution factor}} \times \text{Total plating volume} \quad (1)$$

### Formulation cookies and meatball from *C. indica* L flour and *S. platensis*

At this stage, conducted testing of cookies and meatballs formulations acceptable by the community.

## Result and Discussion

### Proximate Analysis

The data analysis showed that the WRCS flour contained higher carbohydrates than the RRCS counterpart as 67.607% and 57.81% respectively. The WRCS flour contained 8.75% water, 0.29% lipid, 17.41% protein, and 5.94% ash. This content as presented in Table 1.

**Table 1.** Proximate Analysis of white and red rhizome composite flour

Nutrition	RRCS (%)	WRCS (%)
Protein	21.75	17.41
Total Lipid	0.5	0.29
Ash	8.94	5.94
Water	11	8.75
Carbohydrate	57.81	67.61
Fiber	9.99	9.54

The specification of WRCS and RRCS had lowest water content compared to wheat flour Codex 199-1995 which was 14.5% m/m. The results showed the potential



expired date both of composite flour were slower than wheat flour. It is possible to distribute both of composite flour to consumers.

The WRCS and RRCS flour had a total ash content of 5.94% and 8.94%. Ash related to inorganic residue after complete oxidation in food sample in proximate analysis (Ismail, 2017). The major minerals present in food such as Calcium (Ca), Sodium (Na), Potassium (K), Chloride (Cl), Phosphorus (P) and Sulfur (S), while trace mineral is Iodine (I), Zinc (Zn), Selenium (Se), Iron (Fe), Manganese (Mn), Copper (Cu), Cobalt (Co), Molybdenum (Mo), Fluoride (F), Chromium (Cr) and Boron (B). Calcium influences variety of physiological processes to build and maintain bone, as well as normal nerve and muscle function. Phosphorus is a component of adenosine triphosphate (ATP) and nucleic acids. Mg, Co, Se, Zn, Fe, Mn, and Mo are important cofactor enzymes (Gharibzahedi & Jafari, 2017). The ash content was detected at 0.41% in the red rhizome and 0.47% in the white rhizome of *C. indica* L. The *S. platensis*'s ash increase content because it has minerals such as Ca, Fe, P, I, Mg, Zn, Se, Co, Mn, K, and Na. *S. platensis* contains Ca, P, Mg, Fe and Zn respectively 620.8, 790.42, 285.90, 49.5, and 3.95 mg/100 g. (Gutiérrez-Salmeán et al. 2015).

*Protein and amino acids*

Protein is a macronutrient that forms important building blocks in the formation of biomolecules. Increasing protein, dietary fiber, and ash content in composite flour influenced by the addition of *S. platensis*. It was indicated by the elevation in protein from 4.72% to 21.75% and 17.41% in RRCS and WRCS. Protein content in RRCS is the highest from *Abalistes stellaris* (ikan etong) 18,89% and *Euthynnus affinis* (ikan tongkol) 20,27% (Januarita et al., 2021). The protein content found in RRCS composite flour is higher than that found in salted egg yolks, which is 16.29%. The protein content found in RRCS composite flour is higher than that found in salted egg yolks, which is 16.29% (Setyaji & Monica, 2023). This condition is also in the protein content of WRCS composite flour, which is 17,42%. Composite of 10% *S. platensis* on artificial meat with other additions of wheat gluten, 10% soybeans, mushrooms, tofu showed a protein content of 29.32% (Sari et al., 2022).

The changes in dietary fiber content from 0.04% to 9.99% and 9.54%, as well as ash from 1.37% to 8.94% and 5.40%. Plant proteins have important role in regulation of metabolism and modulation to reduce cholesterol, and triglycerides through inhibition of synthase fatty acids which has impact as anti-obesity. Target regulation of cholesterol and triglyceride metabolism through genes that function to reduce cholesterol and triglycerides metabolism as well as dipeptidyl peptidase

IV (DPP-IV). As an antidiabetic effect plant protein works on alpha amylase, alpha glucosidase metabolism gene cause to lower blood sugar (Nagaoka et al. 2021).

Amino acids as a part of protein contain in RRCS and WRCS flour are amino acid essential and non-essential. Essential amino acid contain in both of flour are phenylalanine, valine, threonine, methionine, histidine, leucine, and lysine. Actually, essential amino acids are main role in maintaining tissue and organ functions.

The non-essential amino acids needed for metabolism. Our body can produce or copy it but sometimes the production is insufficient. The types of non-essential found amino acids found in two composite flours are aspartate, serine, glycine, arginine, alanine, proline, and cysteine. Glycine can be synthesized by our body but the production is insufficient (Alves et al., 2019). RRCS and WRCS flour can be as food source containing glycine. Following are the amino acids contained in both type of composite flour compared to *S. platensis* (table 3). Amino acids content of *S. platensis* in both types of composite flour show similarities. Different types of amino acids are tryptophan which only found in *S. platensis* (Bortolini et al., 2022). The absence of tryptophan caused by different types of *S. platensis* culture medium used. RRCS flour also shown to contain higher proline compared to WRCS flour and *S. platensis*. Based on this, it is suspected that proline is not only derived from *S. platensis* but also from red rhizome *C. indica* L. Unlike the case with aspartate which has highest in WRCS, so it is predicted come from white rhizome *C. indica* L. The highest Cysteine found in *S. platensis* and followed by WRCS flour.

**Table 2.** Amino acids in the Composite Flour and Bartolini et al., 2022

Amino acids	RRCS (%)	WRCS (%)	Bartolini (et al., 2022)
Aspartate	0.62	0.75	0.63
Serine	0.31	0.32	0.30
Glutamate	0.82	0.87	0.84
Glycine	0.35	0.37	0.34
Histidine	0.12	0.12	0.11
Arginine	0.49	0.46	0.44
Threonine	0.34	0.37	0.33
Alanine	1.99	0.29	0.50
Proline	1.99	0.29	0.25
Cysteine	0.002	0.03	0.06
Tyrosine	0.18	0.29	0.30
Valine	0.42	0.45	0.42
Methionine	0.08	0.01	0.17
Lysine	0.3	0.33	0.34
Isoleucine	0.34	0.37	0.36
Leucine	0.52	0.59	0.61
Phenylalanine	0.3	0.34	0.33

The amino acid contained in WRCS and RRCS flour were aspartate, serine, alanine, glutamate, glycine, histidine, cysteine, arginine, methionine, glutamate, valine, and isoleucine will be explained as follows. Aspartate in the form of L-aspartate has function as protein biosynthesis and a neurotransmitter (Johnson, 2017). By adding D-aspartate to a neurotransmitter also functions as a regulation in reproduction, hormones, and nerve protection (Li et al., 2018). Serine as a non-essential is a precursor for the biosynthesis of glycine and cysteine, thus the availability of serine is needed for the formation of both types of amino acids.

Alanine serves to avoid the symptoms of paresthesia and reduces urine overflow, while the addition of beta-alanine increases exercises intensity in athletes (Harris & Stellingwerff, 2013). Glutamate is a neurotransmitter to the nerve system, name as presynaptic and postsynaptic (Hao & Plested, 2022). Glutamate can be synthesized from glucose and various nitrogen containing nutrients, as well as glutamine, aspartate, asparagine, serine, glycine, proline, and alanine. On certain circumstances, glutamine, glutamate, glycine, proline, and arginine could not be synthesized in quantity or quality because the influence of pathophysiology.

Imbalance and indispensable amino acids will reduce growth and trigger various diseases. For the example glycine, very effective for promoting growth, improving health, and preventing various diseases including cancer, cardiovascular disease, some inflammatory diseases, obesity, and diabetes. Also influencing sleep quality and neurological function. Also serves as a precursor for several molecular metabolites, such as creatine, glutathione, purines, and porphyrins (Razak et al., 2022).

Histidine include essential amino acids that affect proton buffers, metal ion chelation, scavenging of reactive oxygen and nitrogen species, erythropoiesis, and histaminergic systems. Histidine contain in proteins such as haemoprotein, HIS glycoprotein, hystatins, HIS calcium binding protein and filaggrin, HIS containing dipeptides especially carnosine, methyl and Sulphur. Thus, it contains of HIS (3 methylhistamine, 1 methylhistidine and ergothioneine. HIS as component preservation and myocardial protection in cardiac surgery. HIS supplementation is needed on neurological disorders, atopic dermatitis, metabolic syndrome, diabetes, uremic anemia, ulcers, inflammatory bowel syndrome, malignancies, muscle performance enduring strenuous exercise (Holeček, 2020). Diet methionine 46 mg/ kg/ d, lysine 6.0 g/ d, histidine 8.0 g/ d can reduce fat level and skin condition so can be used for herpes treatment (Cynober et al., 2020). Cysteine functions as an antioxidant has potential to capture Reactive Oxygen Species (ROS). Cysteine function as whitening by

forming pheomelanin which produces yellowish or reddish colors and blocking the formation of eumelanin which produces dark colors (Oshimura & Sakamoto, 2017).

Arginine supplements may increase Growth Hormone (GH) and Thyroid Stimulating Hormone (TSH) (Apolzan et al., 2022). Research shows by giving 10 g to men aged 18 to 39 years increases muscle mass by 18.5 kg / up to 25 kg / m<sup>2</sup> (Liang et al., 2021a). Arginine is required in spermatogenesis of pregnancy and neonates to maintain nitrogen balance (Watford & Wu, 2018). Methionine is a non-essential amino acid precursor cysteine, as well as phenylalanine as a tyrosine precursor. The intake is needed of methionine and phenylalanine in nutrient. A single intake of good leucine supplements with protein supplements can improve of sarcopenia, which is loss of muscle mass because of decreased mobility (Tezze et al., 2023).

Glutamate undertake decarboxylation will transform into Gamma Aminobutyric Acid (GABA) as a neurotransmitter with microbiota's help. Microbiota-gut-brain axis disorder in cases of depression corrected via human microbiota function through transplantation to stimulate GABA (Mayneris-Perxachs et al., 2022).

L-Valine, L-isoleucine, and L-leucine are essential amino acids or synthesized insufficiency in humans and animals. It is referred to as Branched Chain Amino Acids (BCAAs) because they contain branched side chains in scaffolds necessary for growth, especially physiology and biology. Increasement levels of BCAAs in humans will reduce accuracy and development of cardiovascular disease, T2DM metabolic disorders, obesity, and neurodegenerative (Liang et al., 2021b).

#### Lipid and Fatty Acids

The lipid content in RRCS and WRCS flour relatively low as 0.29% and 0.50% compared to standard dietary lipids. People's habits in consuming foods with low lipid levels can minimize the prevalence of obesity (Muriyati et al., 2023) . Both composite flours contain same fatty acid presented in Table 3.

**Table 3.** Fatty acids in the Composite flour of *C. indica L* and *S. platensis*

Fatty Acid	RRCS (%)	WRCS (%)
Myristic Acid	1.29	1.76
Palmitoleic acid	1.60	2.55
Palmitic Acid	38.25	39.79
Linoleic Acid	24.29	25.94
Oleic Acid	21.57	6.13
Stearic Acid	6.13	1.93
Behenic Acid	1.73	0.8
Lignoceric Acid	0	0.8

The highest saturated fatty acids were palmitic as in 39.79% in WRCS and 38.25% in RRCS. Palmitic acid required for average dietary fat intake of 20 g/day for 20-30% of total fatty acids has functions in fetal brain development (Murru et al., 2022). The second largest fatty acid in both flours is linoleic acid (LA) with number in RRCS and WRCS 24,29% - 25,94%. The LA content found in *Terminalia catapa* L seeds shows lower levels compared to those found in RRCS and WRCS, which is 13.09% (Nirwana et al., 2022). In addition to LA also needed intake of Linolene acid (omega 3) which can be found in *Portulaca oleracea* (Dewanti et al., 2023; Muhammad et al., 2023).

LA (omega-6) known as  $\Omega$ -6 polyunsaturated is one of the 6 Polyunsaturated Fatty Acids (PUFA) has essential fatty acids. It affects insulin sensitivity and improves pancreatic  $\beta$ -cell function (Marangoni et al., 2020). LA fatty acid (PUFA) plays a role in glucose homeostasis in regulating blood sugar homeostasis by increasing insulin sensitivity, increasing blood sugar binding due to improved pancreatic beta cell function in sex, genetics and different environmental factors, besides reduce cardiovascular disease (Hamilton & Klett, 2021).

Palmitoleic found in WRCS at 2.55% and 1.6% RRCS flour. Palmitoleic 9-hexadecenoic acid; 16:1 N-7 is a lipokine that functions as a nutraceutical in the treatment of non-alcoholic liver disease. Murinae is inflammation by induction of fat intake of 59% for 12 days, showed increasing insulin tolerance and decreasing serum insulin on day 15 after administration of palmitoleic supplementation 300 mg / kg / day (Souza et al., 2020). Non-essential oleic acid in RRCS flour at 21.57% while WRCS flour is 6.13%. Intake oleic acid can maintain weight via metabolism stimulating AMP activated protein kinase.

In addition, through the prevention of nucleotide binding oligomerization domain like receptor 3/ caspase - 1 inflammasome pathway, induction of oleoylethionamide synthetase and the down regulation stearoyl-Co A desaturated activity (Tutunchi et al., 2020)

Stearic content in RRCS higher than WRCS flour at 6.13% and 1.93%. Stearic acid saturated with 18 carbon long chain fatty acids, also induces apoptosis and autophagy (Yang et al., 2019). Stearic acid can suppress tumor cell growth. Lignoceric acid specifically found in WRCS flour only and has saturated fatty acids. Behenic acid higher in RRCS flour and stable during binding to proteins as ligand complex in tackling T2DM (Nath et al., 2022). Myristic levels almost same in the two composite flours. Myristic is a saturated fatty acid (14:0) that is essential in lipid milks. These fatty acids are beneficial for cardiovascular health. Research shows the presence of myristic acid is associated with increased concentrations of ApoCIII in cardiovascular patients.

Invitro experiments showed myristic acid stimulates ApoCIII in the expression of HepG2 cells and slows down HDL catabolism (Olivieri et al. 2020)

*Fiber*

The composite flour specification shows highest percentage of dietary fiber as 9.99% for WRCS and 9.54% for RRCS flour. Dietary fiber increases health quality by protecting against constipation problems and preventing digestive cancers, regulating blood glucose and decreasing the risk of diabetes, and reducing cardiovascular disease risk by weight loss through lipid regulation. This also suppresses lipid levels by about 5-10% cholesterol, controls blood glucose, improves immune function, and reduces inflammation (Surampudi et al. 2016). Blood glucose reduction influenced by the dietary fiber effect in composite flour.

Fiber consumption 10 g/day for a median intervention duration of 8 weeks in diabetic type 2 patients significantly reduced HbA1c, fasting blood glucose, fasting insulin, and Homeostatic Model Assessment for Insulin Resistance. Fiber foods are effective in improving glycemic control and insulin sensitivity in diabetic type 2 patients (Mao et al., 2021).

*In vitro growth of Lactobacillus*

The total population of *L. casei* Shirota cultured in UMW, M-GCW, and M-GCW-PS showed no significant difference ( $P > 0.05$ ), bacterium was  $8.23 \pm 0.04$ ,  $8.17 \pm 0.02$ , and  $8.17 \pm 0.00$  CFU/mL in the three media. Table 4 shows a significant difference in the total population in M-PSW at  $8.11 \pm 0.03$  CFU/mL. The higher growth in UMW due to immediate fermentation of glucose, while in M-GCW still through the process of starch decomposition into monosaccharides. *L. casei* Shirota cultured on M-PSW requires time to break down proteins to grow into amino acids.

**Table 4.** Total population of *L. casei* Shirota growth in various media

Media	Total Population (CFU/mL)
UMW	$8.23 \pm 0.04$
M-GCW	$8.17 \pm 0.02$
M-PSW	$8.11 \pm 0.03$
M-GCW-PS	$8.17 \pm 0.00$

*S. platensis* contained in the culture media as source of cysteine, proline, glycine and glucose (Zhang et al. 2020). The result of *L. rhamnosus* cultured in UMR and M-GCR-PS showed no significant difference ( $P > 0.05$ ). According to Table 5, the total population in UMR was  $7.79 \pm 0.10$  CFU/mL and M-GCR-PS contained  $7.85 \pm 0.00$  CFU/mL. However, the total  $7.77 \pm 0.08$  CFU/mL and  $7.55 \pm 0.11$  CFU/mL found on the M-PSR and modified M-GCR had a significant difference ( $P < 0.05$ ).



The population of *L. rhamnosus* cultured on media M-GCR lower in numbers. Maybe it is due to higher flavonoid content in the red *C. indica* L rhizome compared to white one. Flavonoids as antibacterials can inhibit nucleic acid synthesis, cytoplasm, membrane permeability, and porin functions, including energy metabolism systems, inhibiting binding and formation of biofilms (Xie et al., 2015). The red rhizome of *C. indica* L contains flavonoids, tannins, and sterols are antimicrobial and anthelmintic (Kumbhar et al., 2018). Rhizome in *C. indica* L also contains alkaloids, saponins, polyphenols and terpenoids (Ifandari et al., 2020). *C. indica* L juga while tannins reduce *Lactobacillus* growth (Peng et al., 2018). Anthocyanins are also mainly antimicrobial *Escherichia coli* dan *Staphylococcus aureus* (Khairi et al., 2023) (Khairi et al., 2023).

The *Lactobacillus* in vitro growth is inhibited by flavonoids and tannins of *C. indica* L, but it can support sterol content (Miedes et al., 2023). Reason for higher growth of *L. rhamnosus* on MRSA media because glucose content can used directly. In modified MRSA, the glucose dominantly source comes from the rhizome of *C. indica* L. Research shows the *C. indica* L rhizome has eccentric hila type of glucose (Huang et al. 2015), while the starch content estimated resistant and has higher concentration than glucose.

**Table 5.** Total Population of *L. rhamnosus* growth in various media

Media	Total Population (CFU/mL)
UMR	7.79 ± 0.10
M-GCR	7.55 ± 0.11
M-PSR	7.77 ± 0.08
M-GCR-PS	7.85 ± 0.00

The content of amino acids, such as methionine, cystine, lysine, and tryptophan, and fatty acids, including palmitic acid, oleic acid, and linoleic acid contained in *S. platensis* supports the growth of *L. rhamnosus*. *S. platensis*. Beta-carotene can be converted into vitamins A and B, as well as minerals by 3-7%.

*C. indica* L contains bioactive hydroxytrimethoprim, 3-7 epoxy-carophyllan 6 one, swietenine, typhasterol, hexacosanedioic acid a 3-beta, 6-alfa, 7-alpha trihydroxy 5 beta choline 24 oleic acid function as antimicrobe, anthelmintic potential and HIV reverse transcriptase inhibition (Kumbhar et al. 2018b). *C. indica* is rich in 67.99% carbohydrate and 22.50% fiber, 319.15 calory/100 grams. Rhyzome contains 24405 ppm potassium, sodium, calcium, and magnesium (7500, 1345 and 1610 ppm) also Cu, Ni, Mn, Fe, Zn (Gaur 2014). *C. Indica* L has potential as functional food, source of vitamins, nutritional and nutraceutical. Biomass and nutrient characteristics are fiber 25.1%, starch 28.5%,

crude oil 4.72% and lipid 5.75% total calories 423 Kcal/100 gr wet weight. Rhizome contains minerals and vitamins. Bioactive compounds such as rosmarinic acid, psoromic acid, usnic acid, isoeugenitol, ellagic acid, coumarin acid and switenine able to protect free radical cells including DNA and protein degradation. Acetone extract in rhizome shows antioxidant activity. Ic 50 values with DPPH, ABTS+, O<sub>2</sub>\_, showing 21.23 and 170 ug/ml. Decreasing of Ferric Reducing Antioxidant Power (FRAP), Cupric Ion Reducing Antioxidant Capacity (CUPRAC) and DNA protection correlated with total phenolids and flavonoids. Beta carotene and linoleic acid show significant acetone extract, approxiamtel IC50 is 2.35 ug/ml and acetone extract IC 50 27.1 ug/ml. *C indica* become healthy potential source of nutrients and metabolism (Ayusman et al., 2020b).

Glucose source of growing culture media of *L. casei* Shirota can be substituted with white rhizome *C. indica* L. Red rhizome has greater potential as glucose substitute of growing of *L. rhamnosus*. *S. platensis* also has potential as a prebiotic and protein source in modified MRSA. As for the two *Lactobacillus*. Based on these facts, white rhizome flour of *C. indica* L. and *S. platensis* has potential of prebiotic system.

*Cookies and meatballs from composite flour*

To make cookies and meatball formulation which are healthy and acceptable by market was going through various tests. The acceptable formulation is using *C. indica* L flour 80 gr, *C. indica* L starch 20 gr, coconut oil 60 gr, coconut sugar 50 gr, milk skim powder 40 gr, egg 40 gr and *S. platensis* powder 1 g. Cookies product result is fibrous, soft, and crunchy. Also has density to ensure it do not crumble easily. Taste close to sweet and has smell of coconut aroma to satisfy consumers. The following is presented a figure describing the product of the research process (Figure 2).



**Figure 2.** Product of research process

Rhizome produced from planting *C. indica* L after being processed into flour is mixed with *S. platensis* which is the result of cultivation. A mixture of the two

types of flour is called composite flour. *Lactobacillus* invitro growth tests showed positive results for two types of composite flour. Furthermore, the composite flour was successfully made into cookies and meatballs that were favored.

The composition flour of *C. indica* and *S. platensis* also using for making meatballs and successfully has savory taste, with chewy texture and delicate meat aroma. Formulation of was diced meat 2000 g, *C. indica* L flour 250 g, sago 250 g, *S. platensis* 1 gram, gelatinous 14 gr, egg 80 g, garlic 100 g, pepper 2 tea spoon, onion 100 g. It acceptable by consumers. Cookies and meatballs are safe to consumer who has gluten allergic because not using wheat flour.

## Conclusion

Based on the consideration of its nutritional content, potential as a prebiotic and can be processed into food variations such as cookies and meatballs, rhizomes, *C. indica* L and *S. platensis* can be used as alternative foods to maintain food security. WRCS and RRCS flour contain carbohydrates, proteins, and lipids that can meet daily nutritional needs and contain prebiotics.

## Acknowledgments

Thank you to the Ministry of Research and Technology and Higher Education of the Republic of Indonesia and to the UAI Institute for Research and Community Service for funding and supporting this research.

## Author Contribution

The idea and conceptual of this research is Nita Noriko. Methodology and formal analysis are Risa Swandari Wijihastuti, Analekta Tiara Perdana and Hidayat Yorianta Sasaerila. Validation data is Alina Primasari.

## Funding

This research won a research funding grant from the Ministry of Research and Technology and Higher Education of the Republic of Indonesia.

## Conflicts of Interest

The results of the research are objective and do not cause problems and conflicts of interest from various parties

## References

- Algar, A. F. C., B.Umali, A., & Tayobong, R. R. P. (2019). Physicochemical and functional properties of starch from philippine edible canna (*canna indica* L.) Rhizomes. *Journal of Microbiology, Biotechnology and Food Sciences*. Retrieved from <https://api.semanticscholar.org/CorpusID:201210880>
- Alves, A., Bassot, A., Bulteau, A.-L., Pirola, L., & Morio, B. (2019). Glycine metabolism and its alterations in obesity and metabolic diseases. *Nutrients*, *11*(6), 1356.
- Amelia, W. R., Leoni, A. P., Syauby, A., Laksmi, P. W., Purba, M. B., Noer, E. R., Wijayanti, A., Sri Nofi, L., Sundari, B., & Arini, M. (2023). Malnutrition at hospital admission and its associated factors in internal medicine inpatients. *Jurnal Gizi Indonesia*, *11*(2), 128-139. <https://doi.org/10.14710/JGI.11.2.128-139>
- Apolzan, J. W., Stein, J. A., Rood, J. C., Beyl, R. A., Yang, S., Greenway, F. L., & Lieberman, H. R. (2022). Effects of acute arginine supplementation on neuroendocrine, metabolic, cardiovascular, and mood outcomes in younger men: A double-blind, placebo-controlled trial. *Nutrition*, *101*, 111658.
- Ayusman, S., Duraivadivel, P., Gowtham, H. G., Sharma, S., & Hariprasad, P. (2020a). Bioactive constituents, vitamin analysis, antioxidant capacity and  $\alpha$ -glucosidase inhibition of *Canna indica* L. rhizome extracts. *Food Bioscience*, *35*, 100544. <https://doi.org/10.1016/j.fbio.2020.100544>
- Ayusman, S., Duraivadivel, P., Gowtham, H. G., Sharma, S., & Hariprasad, P. (2020b). Bioactive constituents, vitamin analysis, antioxidant capacity and  $\alpha$ -glucosidase inhibition of *Canna indica* L. rhizome extracts. *Food Bioscience*, *35*, 100544.
- Bortolini, D. G., Maciel, G. M., Fernandes, I. de A. A., Pedro, A. C., Rubio, F. T. V., Branco, I. G., & Haminiuk, C. W. I. (2022). Functional properties of bioactive compounds from *Spirulina* spp.: Current status and future trends. *Food Chemistry: Molecular Sciences*, *5*, 100134.
- Bvenura, C., & Kambizi, L. (2022). Chapter 5 - Future grain crops. In R. Bhat (Ed.), *Future Foods* (pp. 81-105). Academic Press. <https://doi.org/10.1016/B978-0-323-91001-9.00032-3>
- Campos-Perez, W., & Martinez-Lopez, E. (2021). Effects of short chain fatty acids on metabolic and inflammatory processes in human health. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*, *1866*(5), 158900. <https://doi.org/10.1016/j.bbalip.2021.158900>
- Chouhan, S. (2015). Enumeration and identification of standard plate count bacteria in raw water supplies. *IOSR Journal of Environmental Science*, *9*(2), 67-73. <https://doi.org/10.9790/2402-09226773>
- Cynober, L., Bier, D. M., Stover, P., Kadowaki, M., Morris, S. M., Elango, R., & Smriga, M. (2020). Proposals for Upper Limits of Safe Intake for Methionine, Histidine, and Lysine in Healthy Humans. *The Journal of Nutrition*, *150*, 2606S-2608S. <https://doi.org/10.1093/jn/nxaa231>



- Dewanti, F. D., Sukendah, Lestari Tarigan, P. L., & Yunus, A. (2023). The Effects of various planting media and organic fertilizers on the growth and yield of purslane (*Portulaca oleracea* L.). *Jurnal Penelitian Pendidikan IPA*, 9(6), 4189–4193. <https://doi.org/10.29303/JPPIPA.V9I6.3905>
- Firna, A. L., Bambang, A. N., & Afifah, D. N. (2018). Sensitivitas dan spesifisitas food consumption score dalam mengidentifikasi rawan pangan dengan parameter status gizi balita. *Jurnal Gizi Indonesia*, 6(2), 102–108. <https://doi.org/10.14710/JGI.6.2.102-108>
- Gaur, A. (2014). Nutritive value of rhizome of the *Canna indica* Linn. and characteristics of various extracted materials from the rhizome. *International Journal of Pharma and Bio Sciences*, 5, P644–P648.
- Gharibzahedi, S. M. T., & Jafari, S. M. (2017). The importance of minerals in human nutrition: Bioavailability, food fortification, processing effects and nanoencapsulation. *Trends in Food Science & Technology*, 62, 119–132. <https://doi.org/10.1016/j.tifs.2017.02.017>
- Gutiérrez-Salmeán, G., Fabila-Castillo, L., & Chamorro-Cevallos, G. (2015). Nutritional and Toxicological Aspects of *Spirulina* (*Arthrospira*). *Nutricion Hospitalaria*, 32(1), 34–40. <https://doi.org/10.3305/nh.2015.32.1.9001>
- Hamilton, J. S., & Klett, E. L. (2021). Linoleic acid and the regulation of glucose homeostasis: A review of the evidence. *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 175, 102366. <https://doi.org/10.1016/j.plefa.2021.102366>
- Hao, Y., & Plested, A. J. R. (2022). Seeing glutamate at central synapses. *Journal of Neuroscience Methods*, 375, 109531. <https://doi.org/10.1016/j.jneumeth.2022.109531>
- Harris, R. C., & Stellingwerff, T. (2013). Effect of  $\beta$ -Alanine supplementation on high-intensity exercise performance. In *Nestle Nutrition Institute Workshop Series* (Vol. 76, pp. 61–71). S. Karger AG. <https://doi.org/10.1159/000350258>
- Holeček, M. (2020). Histidine in health and disease: Metabolism, physiological importance, and use as a supplement. *Nutrients*, 12(3), 1–20. <https://doi.org/10.3390/nu12030848>
- Huang, J., Zhao, L., Man, J., Wang, J., Zhou, W., Huai, H., & Wei, C. (2015). Comparison of physicochemical properties of B-type nontraditional starches from different sources. *International Journal of Biological Macromolecules*, 78, 165–172. <https://doi.org/10.1016/j.ijbiomac.2015.04.006>
- Ifandari, Widyarini, S., Hartanto Nugroho, L., & Pratiwi, R. (2020). Phytochemical analysis and cytotoxic activities of two distinct cultivars of ganyong rhizomes (*Canna indica*) against the wide colon cancer cell line. *Biodiversitas*, 21(4), 1660–1669. <https://doi.org/10.13057/biodiv/d210447>
- Indrayani, P. E., Suryani, D., Oktaviyani, P., Suyitno, S., Purwani, S. P. M., & Dharmawan, N. K. S. (2023). The Government's policies on the prevention of Covid-19 transmission impacting household food security in Indonesia. *Jurnal Penelitian Pendidikan IPA*, 9(9), 499–507. <https://doi.org/10.29303/JPPIPA.V9I9.4413>
- Ismail, M. M. S., Piercey-Normore, M. D., & Rampitsch, C. (2018). Proteomic analyses of the cyanobacterium *Arthrospira* (*Spirulina*) *platensis* under iron and salinity stress. *Environmental and Experimental Botany*, 147, 63–74. <https://doi.org/10.1016/j.envexpbot.2017.11.013>
- Ismail, B. P. (2017). Ash Content Determination. In S. S. Nielsen (Ed.), *Food Analysis Laboratory Manual* (pp. 117–119). Springer International Publishing. [https://doi.org/10.1007/978-3-319-44127-6\\_11](https://doi.org/10.1007/978-3-319-44127-6_11)
- Johnson, E. C. (2017). Aspartic Acid☆. In *Reference Module in Biomedical Sciences*. Elsevier. <https://doi.org/10.1016/B978-0-12-801238-3.97338-0>
- Jung, F., Krüger-Genge, A., Waldeck, P., & Küpper, J.-H. (2019a). *Spirulina platensis*, a super food? *Journal of Cellular Biotechnology*, 5, 43–54. <https://doi.org/10.3233/JCB-189012>
- Jung, F., Krüger-Genge, A., Waldeck, P., & Küpper, J.-H. (2019b). *Spirulina platensis*, a super food? *Journal of Cellular Biotechnology*, 5, 43–54. <https://doi.org/10.3233/JCB-189012>
- Kamaludin, A. M. R., & Holik, H. A. (2022). Review article: Chemical content and pharmacological activities of *Spirulina* sp. *Indonesian Journal of Biological Pharmacy*, 2(2), 59–66. Retrieved from <https://journal.unpad.ac.id/ijbp/article/download/38269/pdf>
- Khairi, N., Hapiwaty, S., & Indrisari, M. (2023). Activity, formulation and effectiveness of black rice extract (*Oryza sativa* L) gel against *Staphylococcus aureus* and *Escherichia coli* bacteria. *Jurnal Penelitian Pendidikan IPA*, 9(6), 4273–4278. <https://doi.org/10.29303/JPPIPA.V9I6.3914>
- Khairulbahri, M. (2021). Analyzing the impacts of climate change on rice supply in West Nusa Tenggara, Indonesia. *Heliyon*, 7(12), e08515. <https://doi.org/10.1016/j.heliyon.2021.e08515>
- Kumbhar, S. T., Patil, S. P., & Une, H. D. (2018). Phytochemical analysis of *Canna indica* L. roots and rhizomes extract. *Biochemistry and Biophysics Reports*, 16, 50–55. <https://doi.org/10.1016/j.bbrep.2018.09.002>

- Liang, Y.-F., Long, Z.-X., Zhang, Y.-J., Luo, C.-Y., Yan, L.-T., Gao, W.-Y., & Li, H. (2021a). The chemical mechanisms of the enzymes in the branched-chain amino acids biosynthetic pathway and their applications. *Biochimie*, 184, 72–87. <https://doi.org/10.1016/j.biochi.2021.02.008>
- Liang, Y.-F., Long, Z.-X., Zhang, Y.-J., Luo, C.-Y., Yan, L.-T., Gao, W.-Y., & Li, H. (2021b). The chemical mechanisms of the enzymes in the branched-chain amino acids biosynthetic pathway and their applications. *Biochimie*, 184, 72–87. <https://doi.org/10.1016/j.biochi.2021.02.008>
- Li, Y., Han, H., Yin, J., Li, T., & Yin, Y. (2018). Role of D-aspartate on biosynthesis, racemization, and potential functions: A mini-review. *Animal Nutrition*, 4(3), 311–315. <https://doi.org/10.1016/j.aninu.2018.04.003>
- Mao, T., Huang, F., Zhu, X., Wei, D., & Chen, L. (2021). Effects of dietary fiber on glycemic control and insulin sensitivity in patients with type 2 diabetes: A systematic review and meta-analysis. *Journal of Functional Foods*, 82, 104500. <https://doi.org/10.1016/j.jff.2021.104500>
- Marangoni, F., Agostoni, C., Borghi, C., Catapano, A. L., Cena, H., Ghiselli, A., La Vecchia, C., Lercker, G., Manzato, E., Pirillo, A., Riccardi, G., Risé, P., Visioli, F., & Poli, A. (2020). Dietary linoleic acid and human health: Focus on cardiovascular and cardiometabolic effects. In *Atherosclerosis* (Vol. 292, pp. 90–98). Elsevier Ireland Ltd. <https://doi.org/10.1016/j.atherosclerosis.2019.11.018>
- Mayneris-Perxachs, J., Castells-Nobau, A., Arnoriaga-Rodríguez, M., Martín, M., de la Vega-Correa, L., Zapata, C., Burokas, A., Blasco, G., Coll, C., Escrichs, A., Biarnés, C., Moreno-Navarrete, J. M., Puig, J., Garre-Olmo, J., Ramos, R., Pedraza, S., Brugada, R., Vilanova, J. C., Serena, J., ... Fernández-Real, J. M. (2022). Microbiota alterations in proline metabolism impact depression. *Cell Metabolism*, 34(5), 681–701.e10. <https://doi.org/10.1016/j.cmet.2022.04.001>
- Miedes, D., Makran, M., Cilla, A., Barberá, R., Garcia-Llatas, G., & Alegría, A. (2023). Aging-related gastrointestinal conditions decrease the bioaccessibility of plant sterols in enriched wholemeal rye bread: in vitro static digestion. *Food & Function*.
- Muhammad, N., Abdullatif, Z., Samad, S., & DAS, S. S. (2023). Analysis of omega-3 content in different cultivation treatments of purslane plants (*Portulaca oleracea*). *Jurnal Penelitian Pendidikan IPA*, 9(9), 7079–7088. <https://doi.org/10.29303/JPPPIPA.V9I9.4423>
- Muliarta, I. N., Hariani, C. Y., Wahyuni, N. M. I., & Rismayanti, A. A. T. (2023). Study of potential food waste in zero waste cities area- Saridewi, Denpasar-Bali. *Jurnal Penelitian Pendidikan IPA*, 9(5), 2595–2603. <https://doi.org/10.29303/JPPPIPA.V9I5.3019>
- Muriyati, M., Hamdana, H., Asri, A., Safruddin, S., & Asnidar. (2023). Fat and carbohydrates as causative factors of obesity of youths at Bulukumba city, South Sulawesi. *Jurnal Penelitian Pendidikan IPA*, 9(5), 2726–2731. <https://doi.org/10.29303/JPPPIPA.V9I5.3467>
- Murru, E., Manca, C., Carta, G., & Banni, S. (2022). Impact of dietary palmitic acid on lipid metabolism. *Frontiers in Nutrition*, 9, 861664. <https://doi.org/10.3389/fnut.2022.861664>
- Nagaoka, S., Takeuchi, A., & Banno, A. (2021). Plant-derived peptides improving lipid and glucose metabolism. *Peptides*, 142, 170577. <https://doi.org/10.1016/j.peptides.2021.170577>
- Nath, V., Paul, R. K., Kumar, N., & Kumar, V. (2022). Identification of behenic acid as medicinal food for the diabetes mellitus: structure-based computational approach and molecular dynamics simulation studies. *Journal of Molecular Modeling*, 28(3), 73. <https://doi.org/10.1007/s00894-022-05060-8>
- Nirmalawaty, A., Panjaitan, T. W. S., & Alamsyah, A. (2023). Preference responses of wheat flour substitution with jackfruit seed flour on oyster mushroom nuggets. *Jurnal Penelitian Pendidikan IPA*, 9(7), 5753–5758. <https://doi.org/10.29303/JPPPIPA.V9I7.4672>
- Nirwana, A. R., Suhendra, D., & Gunawan, E. R. (2022). Separation of n-methylhydroxamic fatty acids based on ketapang seed oil using high performance liquid chromatography. *Jurnal Penelitian Pendidikan IPA*, 8(5), 2329–2334. <https://doi.org/10.29303/JPPPIPA.V8I5.1611>
- Olivieri, O., Speziali, G., Castagna, A., Pattini, P., Udali, S., Pizzolo, F., Liesinger, L., Gindlhuber, J., Tomin, T., Schittmayer, M., Birner-Gruenberger, R., Cecconi, D., Girelli, D., Friso, S., & Martinelli, N. (2020). The Positive Association between Plasma Myristic Acid and ApoCIII Concentrations in Cardiovascular Disease Patients Is Supported by the Effects of Myristic Acid in HepG2 Cells. *The Journal of Nutrition*, 150(10), 2707–2715. <https://doi.org/10.1093/jn/nxaa202>
- Oshimura, E., & Sakamoto, K. (2017). Amino acids, peptides, and proteins. In K. Sakamoto, R. Y. Lochhead, H. I. Maibach, & Y. Yamashita (Eds.), *Cosmetic Science and Technology* (pp. 285–303). Elsevier. <https://doi.org/10.1016/B978-0-12-802005-0.00019-7>

- Peng, K., Jin, L., Niu, Y. D., Huang, Q., McAllister, T. A., Yang, H. E., Denise, H., Xu, Z., Acharya, S., Wang, S., & Wang, Y. (2018). Condensed tannins affect bacterial and fungal microbiomes and mycotoxin production during ensiling and upon aerobic exposure. *Applied and Environmental Microbiology*, 84(5), 2274–2291. [https://doi.org/10.1128/AEM.02274-17/SUPPL\\_FILE/ZAM005188336S1.PDF](https://doi.org/10.1128/AEM.02274-17/SUPPL_FILE/ZAM005188336S1.PDF)
- Prijanti, A., Nurhayati, R., Iswanti, F., Khoiriyah, Z., Paramita, R., & Fadilah, F. (2022). Java Sea *Spirulina platensis* chemical analysis and its protective ability against H<sub>2</sub>O<sub>2</sub>-exposed umbilical cord mesenchymal stem cells according to CD73, CD90, and CD105, viability, and HIF-1 alpha docking. *Journal of Applied Pharmaceutical Science*, 12, 67–75. <https://doi.org/10.7324/JAPS.2022.120908>
- Razak, M. A., Begum, P. S., Viswanath, B., & Rajagopal, S. (2022). Erratum: Multifarious Beneficial Effect of Nonessential Amino Acid, Glycine: A Review (Oxidative Medicine and Cellular Longevity (2017) 2017 (1716701) DOI: 10.1155/2017/1716701). In *Oxidative Medicine and Cellular Longevity* (Vol. 2022). <https://doi.org/10.1155/2022/9857645>
- Riandini, E., Susanto, E., Febrianti, N. F., Biologi, J., Matematika, F., & Ilmu, D. (2023). Diversity of canna on Enggano Island, Bengkulu Province. *Jurnal Biologi Tropis*, 23(4), 179–184. <https://doi.org/10.29303/JBT.V23I4.5523>
- Saharan, V., & Jood., S. (2017). NUTRITIONAL COMPOSITION OF SPIRULINA PLATENSIS POWDER AND ITS ACCEPTABILITY IN FOOD PRODUCTS. *International Journal of Advanced Research*, 5(6), 2295–2300. <https://doi.org/10.21474/IJAR01/4671>
- Sari, B. L., Dewi, E. N., & Fahmi, A. S. (2022). Pengaruh penambahan spirulina platensis sebagai sumber protein nabati pada daging analog bagi vegetarian. *Jurnal Mutu Pangan: Indonesian Journal of Food Quality*, 9(2), 76–83. <https://doi.org/10.29244/JMPI.2022.9.2.76>
- Sasaerila, Y. H., Sakinah, S., Noriko, N., & Wijihastuti, R. S. (2021). Effects of Light Environments on Leaf Traits and Phenotypic Plasticity of *Canna indica*. *Biosaintifika: Journal of Biology & Biology Education*, 13(2), 169–177. <https://doi.org/10.15294/biosaintifika.v13i2.30175>
- Setyaji, H., & Monica, M. (2023). Proximate quality of low-sodium salted egg. *Jurnal Penelitian Pendidikan IPA*, 9(10), 7944–7949. <https://doi.org/10.29303/JPPIPA.V9I10.5293>
- Silva, Y. P., Bernardi, A., & Frozza, R. L. (2020). The Role of Short-Chain Fatty Acids From Gut Microbiota in Gut-Brain Communication. In *Frontiers in Endocrinology* (Vol. 11). <https://doi.org/10.3389/fendo.2020.00025>
- Souza, C. O., Teixeira, A. A. S., Biondo, L. A., Silveira, L. S., de Souza Breda, C. N., Braga, T. T., Camara, N. O. S., Belchior, T., Festuccia, W. T., Diniz, T. A., Ferreira, G. M., Hirata, M. H., Chaves-Filho, A. B., Yoshinaga, M. Y., Miyamoto, S., Calder, P. C., Sethi, J. K., & Rosa Neto, J. C. (2020). Palmitoleic acid reduces high fat diet-induced liver inflammation by promoting PPAR- $\gamma$ -independent M2a polarization of myeloid cells. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*, 1865(10), 158776. <https://doi.org/10.1016/j.bbailip.2020.158776>
- Suhaimi, A., Roni, M., & Kusumayana, P. (2023). Analysis of diversity of food consumption in families of stunting toddlers. *Jurnal Penelitian Pendidikan IPA*, 9(10), 8251–8257. <https://doi.org/10.29303/JPPIPA.V9I10.5575>
- Surampudi, P., Enkhmaa, B., Anuurad, E., & Berglund, L. (2016). Lipid Lowering with Soluble Dietary Fiber. *Current Atherosclerosis Reports*, 18(12), 75. <https://doi.org/10.1007/s11883-016-0624-z>
- Syakirotin, M., Karyani, T., & Noor, T. I. (2022). Ketahanan pangan sebelum dan selama pandemi Covid-19 di Kabupaten Bandung. *Jurnal Ilmu Pertanian Indonesia*, 27(3), 472–490. Retrieved from <https://journal.ipb.ac.id/index.php/JIPI/article/view/40524>
- Tezze, C., Sandri, M., & Tessari, P. (2023). Anabolic Resistance in the Pathogenesis of Sarcopenia in the Elderly: Role of Nutrition and Exercise in Young and Old People. *Nutrients*, 15(18), 4073.
- Tutunchi, H., Ostadrahimi, A., & Saghafi-Asl, M. (2020). The Effects of Diets Enriched in Monounsaturated Oleic Acid on the Management and Prevention of Obesity: a Systematic Review of Human Intervention Studies. *Advances in Nutrition*, 11(4), 864–877. <https://doi.org/10.1093/advances/nmaa013>
- Watford, M., & Wu, G. (2018). Protein. *Advances in Nutrition*, 9(5), 651–653. <https://doi.org/10.1093/advances/nmy027>
- Widawati, D., Santosa, G. W., & Yudiati, E. (2022). Pengaruh pertumbuhan spirulina platensis terhadap kandungan pigmen beda salinitas. *Journal of Marine Research*, 11(1), 61–70. <https://doi.org/10.14710/JMR.V11I1.30096>
- Wing, I. S., De Cian, E., & Mistry, M. N. (2021). Global vulnerability of crop yields to climate change. *Journal of Environmental Economics and Management*, 109, 102462. <https://doi.org/10.1016/j.jjeem.2021.102462>
- WU, J., ZHANG, J., GE, Z., XING, L., HAN, S., SHEN, C., & KONG, F. (2021). Impact of climate change on



- maize yield in China from 1979 to 2016. *Journal of Integrative Agriculture*, 20(1), 289-299. [https://doi.org/10.1016/S2095-3119\(20\)63244-0](https://doi.org/10.1016/S2095-3119(20)63244-0)
- Wulandari, E. Y. (2023). Angka lempeng total, most probable number, dan identifikasi bakteri Coliform pada susu sapi segar di Kabupaten Banyuwangi. *Journal of Indonesian Medical Laboratory and Science*, 4(1), 45-58. <https://doi.org/10.53699/JOIMEDLABS.V4I1.114>
- Xie, Y., Yang, W., Tang, F., Chen, X., & Ren, L. (2015). Antibacterial activities of flavonoids: structure-activity relationship and mechanism. *Current Medicinal Chemistry*, 22(1), 132-149. <https://doi.org/10.2174/0929867321666140916113443>
- Yang, Y., Huang, J., Li, J., Yang, H., & Yin, Y. (2019). Effects of Stearic Acid on Proliferation, Differentiation, Apoptosis, and Autophagy in Porcine Intestinal Epithelial Cells. *Current Molecular Medicine*, 20(2). <https://doi.org/10.2174/1566524019666190917144127>
- Zhang, Y., Meng, L., Ai, M., Qiao, Y., Liu, G., Fan, X., Lv, X., & Feng, Z. (2020). Nutrient requirements of *Lactobacillus casei* Shirota and their application in fermented milk. *LWT*, 118, 108735. <https://doi.org/10.1016/j.lwt.2019.108735>